

NEAR EAST UNIVERSITY
GRADUATE SCHOOL OF SOCIAL SCIENCES
APPLIED (CLINICAL) PSYCHOLOGY MASTER PROGRAM



MASTER THESIS

**THE PREVALENCE AND RISK FACTORS OF
CIGARETTE AND ALCOHOL USE AND THEIR
EFFECT ON OTHER PSYCHOACTIVE
SUBSTANCE USE IN TURKISH REPUBLIC OF
NORTHERN CYPRUS**

ELİF CEREN SERT

20131799

THESIS SUPERVISOR

PROF. DR. MEHMET ÇAKICI

NICOSIA, 2015

NEAR EAST UNIVERSITY
GRADUATE SCHOOL OF SOCIAL SCIENCES
APPLIED (CLINICAL) PSYCHOLOGY
POSTGRADUATE PROGRAM



MASTER THESIS
THE PREVALENCE AND RISK FACTORS OF CIGARETTE AND
ALCOHOL USE AND THEIR EFFECT ON OTHER PSYCHOACTIVE
SUBSTANCE USE IN TURKISH REPUBLIC OF NORTHERN CYPRUS

Prepared by; Elif Ceren SERT

Examining Committee in Charge

Assist. Prof. Dr. Zihniye OKRAY

Chairman of the Committee

Department of Psychology

European University of

Lefke

Prof. Dr. Mehmet ÇAKICI

Department of Psychology

Near East University

(Supervisor)

Assist. Prof. Dr. Deniz KARADEMİR

Department of Psychology

Near East University

Approval of the Graduate School of Social Sciences

Prof. Dr. Çelik Aruoba- Dr. Muhittin Özsağlam

ÖZET

KUZHEY KIBRIS TRK CUMHURİYETİNDE SİĞARA VE ALKOL KULLANIMININ YAYGINLIĞI, RİSK FAKTÖRLERİ VE DİĞER PSİKOAKTİF MADDE KULLANIMINA ETKİSİ

Hazırlayan: Elif Ceren SERT

Eyll, 2015

Bağımlılık yapıcı madde kullanımı tm dnyada olduđu gibi Kıbrısta danemli halk sağığı sorunlarından biridir. Bu tez alıřması, sigara ve alkol kullanımının yaygınlığının araştırılması, bu sreteki risk faktrlerinin belirlenmesi ve buradan hareketle diğerk psikoaktif madde kullanımına olan etkinin tespitini amalamaktadır. alıřmanın evreni Kuzey Kıbrıs'ta 18-65 yař arasında KKTC'de yařayan ve Trke konuşan bireylerden oluřmaktadır. Arařtırmanın rneklem grubunu Kıbrısta yařayan kotalı ok basamaklı tabakalandırılmıř sekisiz 994 kiři oluřturmuřtur. alıřmanın anket formu akıcı ve ark.'nın (2003) Trke'ye uyarladığı Avrupa Konseyi'nin "Model Avrupa Anketi" (The Model European Questionnaire) isimli anket alıřmasından yararlanılarak hazırlanmıřtır. KKTC'de yetiřkinler arasında yařam boyu sigara ime oranı %62,7, alkol kullanma yaygınlığı %77,1 ve yasa dıřı madde kullanma oranı %7,7 olarak tespit edilmiřtir. řimdiki sigara kullanma oranı ise %41,8 olarak tespit edilmiřtir. Sigara kullananların %4,3' 11 yař altında, %57,6'sı ise 18 ve stnde sigara kullanmaya bařladıklarını belirtmiřlerdir. Alkol kullananların ise 11 %1,5 yař altında, %62,8 ise 18 yař ve stnde kullanmaya bařlamıřlardır. Bir defada 5 bardaktan fazla ienler %8,3'tr. Alkoll ikiler daha fazla eğıence, arkadaşlarla, rahatlama ve stres atmak iin kullanıldığı grlmektedir. Erkekler kadınlardan daha ok alkol kullandığı grlmektedir. Her gelirden insanın alkol itiğı grlmektedir. Alkol kullanmayanların dine daha ok nem verdikleri grlmektedir. alıřmada sigara ve alkol kullananlarda kullanmayanlara gre daha ok DPM ve yasadıřı madde kullanımı grlmřtr. Arařtırma sonularına bakıldığında sigara ve alkol kullanımının KKTC'de yaygın olarak kullanıldığını ve yasadıřı psikoaktif maddeler iin bir risk faktr olduklarını ortaya koymaktadır. Bu iliřkiden yola ıkarak DPM kullanımını nlemek iin sigara

ve alkol kullanımına yönelik önleyici eğitim programlarına ihtiyaç bulunmaktadır. Sigara ve alkol kullanımına yönelik ilkokul çağından itibaren yapılacak eğitim ve önleme çalışmaları hem sigara ve alkol kullanımındaki yaygınlığı azaltacak hem de DPM kullanımını da aynı zamanda azaltacaktır. Sonuç olarak sigara ve alkole yönelik KKTC’de multidisipliner yaklaşımla bir halk sağlığı politikasına ihtiyaç bulunmaktadır.

THE PREVALENCE AND RISK FACTORS OF CIGARETTE AND ALCOHOL USE AND THEIR EFFECT ON OTHER PSYCHOACTIVE SUBSTANCE USE IN TURKISH REPUBLIC OF NORTH CYPRUS

Prepared by: Elif Ceren SERT

September, 2015

ABSTRACT

Psychoactive substance use is a very serious public health problem in TRNC as in the world. This research aims to examine the prevalence of cigarette and alcohol, determine the risk factors and its effect on other psychoactive substances (OPS). The setting of the study is in TRNC and the study includes individuals aged between 18 and 65 living in TRNC and speaking Turkish. Sample group in the study is composed of 994 participants in TRNC based on quota multiple-stage randomized sample. The survey used in the study was prepared based on the "The Model European Questionnaire" transcribed by Çakıcı et al. (2003). The lifetime prevalence of cigarette use among adults in TRNC was found to be 62.7%, alcohol use was 77.1% and illicit drug use was 7.7%. The current rate of smoking is 41.8%. 4.3% of smokers stated that they started smoking under the age of 11, whereas 57.6% of them indicated that they started smoking at 18 and above. However, 1.5% of individuals drinking alcohol stated that they started drinking at the age of 11, while 62.8 of them stated that they started drinking at 18 and above. The number of individuals drinking more than five glasses at once is 8.3%. Alcoholic drinks are usually consumed for having fun with friends, relaxing and destressing. It was found that males consumed more alcohol than females. It was also found that individuals with different incomes drank alcohol. It was found that non-drinkers attached more importance to religion. It was further found that drinkers and smokers showed more tendency to use OPS and illegal substance. The results of the study show that smoking and drinking alcohol is high in TRNC and that both smoking and drinking pose a risk factor for illegal psychoactive substances. Based on this relationship, tobacco and alcohol prevention programs are required for preventing OPS use. Prevention programs and prevention

education need to be started from primary school age will decrease the use of alcohol, tobacco and OPS. As a result, public health policy with multidisciplinary approach to smoking and drinking is needed in TRNC.

ACKNOWLEDGEMENT

First, I would like to thank my supervisor Prof. Dr. Mehmet akıcı for her advices and useful directions that make me work on this dissertation more effectively and motivated, I am grateful he was my supervisor. I would like to thank to Assoc. Prof. Dr. Ebru Tansel akıcı, for hersupport, suggestions and guidance during my study. Her evaluations guided me in anacademic sense. I would like to thank my dear teachers Dr. Deniz Ergün, Assoc. Prof. Dr. Ülgen H. Okyayuz, Assist. Prof. Dr. Zihniye Okray and Assist. Prof. Dr. İrem Erdem Atak for the support, understanding, helping and motivation that they provided me during my whole master educations and university. I would also like to thank to Uzm. Psk. Meryem Karaaziz. She has been answering my questions and for helping.

Lastly, I would like to express special gratitude to my family. My mother Nilgün Sert and my father İlyas Sert and my sister Selen Sert have been very supportive, helpful and patient to me in this process. I would also like to my friends for helping and each timeanswering my questions sincerely and guide me in thesis process.

Elif Ceren Sert, Nicosia,

September, 2015

CONTENTS

THESIS APPROVAL PAGE.....	I
ÖZET.....	IV
ABSTRACT.....	VI
ACKNOWLEDGEMENT.....	VIII
CONTENTS.....	IX
LIST OF TABLES.....	XI
ABBREVIATIONS.....	XV
1. INTRODUCTION.....	1
1.1. General Evaluation.....	1
1.2. Cigarette.....	2
1.2.1. Definition of Cigarette.....	2
1.2.2. History of Cigarette.....	2
1.2.3. Prevalence of Cigarette.....	2
1.2.4. Why People Smoke.....	4
1.2.4.1. Biological Reasons.....	4
1.2.4.2. Psychological Reasons.....	4
1.2.4.3. Familial Reasons.....	5
1.2.5. Harms of Cigarette Smoking.....	5
1.2.6. Treatment of Smoking Cessation.....	6
1.3. ALCOHOL.....	8
1.3.1. General Information About Alcohol.....	8
1.3.2. Use of Alcohol.....	8
1.3.3. Alcohol Abuse.....	9
1.3.4. History of Alcohol.....	10
1.3.5. Prevalence of Drinking Alcohol.....	11
1.3.6. Reasons for Drinking Alcohol.....	12
1.3.7. Harms of Drinking Alcohol.....	12
1.3.8. Treatment for Alcohol Problems.....	12
1.4. Psychoactive substance use and reasons.....	15
1.4.1. Psychological Reasons.....	15
1.4.2. Genetic Causes.....	15
1.4.3. Biological Causes.....	15
1.4.4. Sociocultural Causes.....	16
1.4.5. Frequency of Psychoactive Substance Use.....	16

2. METHOD	18
2.1. Sample	18
2.2. Survey	18
2.2.1. The Model European Questionnaire	18
2.3. Procedure	19
2.4. Statistical Method	20
3. RESULT	21
4. DISCUSSION	54
4.1. Cigarette Use	54
4.2. Alcohol Use	55
5. CONCLUSION	57
6. REFERENCES	58
7. APPENDIXES	65
7.1. Appendix 1. Informed Consent Form	65
7.2. Appendix 2. Demographic Information Form	67
The Model European Survey	67
8. CURRICULUM VITAE	92

LIST OF TABLES

Table 1. Comparison of male and female distribution of participants in terms of smoking cigarette, pipe or cigar.....	21
Table 2. Comparison of male and female distribution of the time of serving the highest amount of alcohol in participant houses	22
Table 3. Comparison of male and female age distribution of participants to start smoke	23
Table 4. Comparison of male and female age distribution of participants in terms of drinking constantly more than their peers.....	24
Table 5. Comparison of male and female age distribution of participants in terms of beginning to drink.....	25
Table 6. Comparison of male and female distribution of life long use of smoking.....	26
Table 7. Comparison of male and female distribution of smoking in the past 12 months.....	26
Table 8. Comparison of male and female distribution of smoking in the past 30 days	27
Table 9. Comparison of participants in terms of whether they faced difficulty in quitting smoking or not	28
Table 10. Comparison of participants in terms of consuming alcoholic drinks in the past 12 months	28
Table 11. Comparison of participants in terms of consuming alcoholic drinks in the past 30days.....	29
Table 12. Comparison of participants in terms of consuming alcohol in the past 30 days.....	30
Table 13. Comparison of the amount of alcohol consumed by participants in one go (one drink, a bottle or a glass of beer, a glass of champagne, a glass of raki or other alcoholic drinks).....	30
Table 14. Comparison of last places where participants are being drun.....	31
Table 15. Comparison of participants in terms of how many times they got drunk in their lives.....	32
Table 16. Comparison of participants being drunk in the past 12 months.....	32
Table 17. Comparison of participants being drunk in the past 30 days.....	33
Table 18. Comparison of reasons why participants got drunk.....	34

Table 19. Comparison of gender distribution of smokers and non-smokers.....	35
Table 20. Comparison to place of birth distribution of smokers and non smokers.....	35
Table 21. Comparison to age distribution of smokers and non-smokers	36
Table 22. Comparison of smokers and non-smokers in terms of who they live with.....	36
Table 23. Comparison of education level distribution of smokers and non-smokers.....	37
Table 24. Comparison of education level distribution to fathers of smokers and non-smokers.....	37
Table 25. Comparison of education level distribution to mothers of smokers and non-smokers.....	38
Table 26. Comparison to birth place distribution to mothers of smokers and non-smokers.....	38
Table 27. Comparison to birth place distribution of fathers to smokers and non-smokers.....	39
Table 28. Comparison of distribution to places where smokers and non-smokers live.....	39
Table 29. Comparison of work status distribution of smokers and non-smokers.....	40
Table 30. Comparison of social support distribution of smokers and non-smokers.....	40
Table 31. Comparison of income status distribution of smokers and non-smokers.....	40
Table 32. Comparison of smokers and non-smokers in terms of where they receive most information about smoking.....	41
Table 33. Comparison of smokers and non-smokers in terms of the importance of religion in their lives.....	41
Table 34. Comparison of gender distribution of drinkers and non-drinkers.....	42
Table 35. Comparison of birthday distribution of drinkers and non-drinkers.....	42
Table 36. Comparison of distribution of drinkers and non-drinkers in terms of who they live with.....	43

Table 37. Comparison of education level distribution of drinkers and non-drinkers.....	43
Table 38. Comparison of education level distribution of fathers to drinkers and non-drinkers	44
Table 39. Comparison of education level distribution of mothers to drinkers and non-drinkers.....	44
Table 40. Comparison of birth place distribution of drinkers and non-drinkers.....	45
Table 41. Comparison of birth place distribution to mothers to drinkers and non-drinkers.....	45
Table 42. Comparison of birth place distribution of fathers to drinkers and non-drinkers.....	46
Table 43. Comparison of distribution to places where drinkers and non-drinkers live.....	46
Table 44. Comparison of work status distribution of drinkers and non-drinkers.....	46
Table 45. Comparison of income status distribution to drinkers and non-drinkers.....	47
Table 46. Comparison of drinkers and non-drinkers in terms of where they receive most information about drinking.....	47
Table 47. Comparison of drinkers and non-drinkers in terms to the importance of religion in their lives.....	48
Table 48. Comparison of smokers and non-smokers in terms of OPS use.....	48
Table 49. Comparison of smokers and non-smokers in terms of illegal OPS use.....	49
Table 50. Comparison of smokers and non-smokers in terms of drinking age.....	49
Table 51. Comparison of smokers and non-smokers in terms of constant drinking age compared to their peers.....	49
Table 52. Comparison of smokers and non-smokers in terms of the amount of alcohol consumed in one go.....	50
Table 53. Comparison of smokers and non-smokers in terms of prevalence of drunkenness.....	50
Table 54. Comparison of drinkers and non-drinkers in terms of OPS use.....	50

Table 55. Comparison of drinkers and non-drinkers in terms of illegal OPS use.....	51
Table 56. Comparison of drinkers and non-drinkers in terms of smoking cigarette, pipe or cigar.....	51
Table 57. Odds ratio and confidence intervals of demographic variables for formation of tobacco use obtained from multivariate logistic regression.....	52
Table 58. Odds ratio and confidence intervals of demographic variables for formation of alcohol use obtained from multivariate logistic regression.....	53

ABBREVIATIONS

TRNC: Turkish Republic Northern Cyprus

T.V: Television

U.S: United States

WHO: World Health Organization

OPS: Other Psychoactive Substance

1. INTRODUCTION

1.1. General Evaluation

Nowadays, psychoactive substance use has become a global social problem affecting the whole world. In recent years, struggle against psychoactive substance use has become the most significant agenda in most countries. Psychoactive substance use affects health and leads to death. Psychoactive substance affects lives of people as well as being harmful to health (Turhan, 2011, 33).

It is stated that drugs such as chemical psychoactive substance cause physical and psychological addiction and are used lifelong (Ögel, 1997, 54). When psychoactive substances which may lead to addiction after being used several times are used as a means of trade, they may be the most significant source of income for who want to generate income illegally (Derdiman, 2006, 103). It is stated that illegal psychoactive substances contribute to black economy significantly. It is seen that psychoactive substance use has been increasing significantly especially after World War II and become one of the most significant problems in the world (Köknel, 1998). Cigarette and alcohol are the most widely used psychoactive substances. It is seen that psychoactive substance use usually starts with smoking and continues with the use of other psychoactive substances in time (Tanrıku et al, 2008, 101). In a study by McKee and his friends in Canada, it was stated that 74% of students both smoke and drink alcohol (McKee et al, 2004, 111). In another study it was shown that drinking alcohol increases smoking significantly among students (Keskinöglü et al, 2006, 190). The effect of smoking on drinking alcohol was proved and in literature data on the effect of volatile psychoactive substance and other drugs on the increasing use of these substances exist (McKee et al, 2004, 112). As understood, some addictions trigger each other and lead to other problem addictions (Esirgemez, 2014, 41). Thus, it can be said that the process which starts with smoking, continues with drinking alcohol and ends with using psychoactive substance. It is also stated that using tobacco and drinking alcohol are considered as risk factors for drug use (Esirgemez, 2014, 41).

1.2. Cigarette

1.2.1. Definition of Cigarette

Cigarette is made from the leaves of tobacco and formed from rolling papers around tobacco. It is stated that cigarette contains 400 toxic psychoactive substances (Ögel, 1997, 54). Cigarette is a kind of thin layered tobacco rolled in a paper and practical to carry and use. Tobacco is on the top of the addictive psychoactive substances that are widely used all over the world. It is stated that 70% of people who quit using tobacco are likely to restart using it at the end of the first month. This percentage shows the striking effect of tobacco on becoming addiction (Ögel, 1997, 56).

1.2.2. History of Cigarette

Use of tobacco, which is the oldest and most common habit in the modern era, is one of the biggest social problems. It can be said that use of tobacco becomes more popular every day by taking hold of young people. Cigarette is prepared from the dried leaves of tobacco. As well as being used as a cigar or by chewing, tobacco can also be used with a pipe (Mangır et al, 1992, 17).

America is the homeland of tobacco production and Christopher Colombu is the first person to introduce tobacco to Europe. Tobacco was presented to the queen in Paris by the French ambassador, Jean Nicot, in Portugal in 1600 and planted in the garden of the palace. The most poisonous psychoactive substance in tobacco, nicotine is named after the ambassador (Bariş and İzzettin, 1994, 16). It is stated that while age of drug use and alcohol decreases day by day in Turkey, age of smoking has decreased to age 7.

1.2.3. Prevalence of Cigarette Use

World Health Organization (WHO, 2004) noted that smoking causes the death of 5 million people every year and the number is expected to double in the next 20 years. According to the estimates made by WHO, whereas today the number of smokers is around 1.3 billion, it will expected rise up to 1.7 billion in 2025. Every 5-

5 second a person dies of smoking in the world and this situation causes \$200 million damage to the world economy (DDK, Araştırma ve İnceleme Raporu, 48).

Turkey is one of the countries where cigarette is produced and consumed most. For this reason, smoking has been identified with Turkish identity over time. This caused smoking to become a traditional consumable psychoactive substance. In the last decade, cigarette production increased by 50% in Turkey, which is seen as a serious increase (Bilir et al, 2007, 22). In another study carried out by Ulukoca, Gökgez and Karakoç (2013), it was found that 45.4% of young people smoke and the rate of those smoking regularly every day is 34.9%. In 2008, Turkey ranked the second country consuming cigarette after Pakistan with this tragic increase. Research shows that 750 thousand adolescents start smoking in Turkey every year (Sezer, 1984, 11). The annual rate of expense in smoking increased to \$6.5 billion in Turkey, where the number of smokers is predicted to be 17 million. According to the scientific studies, while 5 million people die of smoking in the world every year, 100.000 people die of smoking in Turkey every year. It is also stated that Turkey has 3 million people with chronic lung disease, 4 million people with asthma and around 50 thousand people are diagnosed with lung cancer every year in Turkey.

In recent years studies that measure tobacco use prevalence is increased in TRNC. First study is conducted at 1996 among high school students and covered 2215 participants and at least once in their life time tobacco use rate is found as % 42 (Çakıcı M & Çakıcı E, 2000a). In the four subsequent high school studies; % 40.6 at 1999, %35.2 at 2004, %26.8 at 2011 and %31 at 2015 (Çakıcı M & Çakıcı E, 2000b, Çakıcı et al 2010, Eş 2015, Çakıcı et al, 2015). The study, which is conducted by Çakıcı et al. (2014) aims to examine the prevalence and risks of psychoactive substance use among university students in TRNC, shows that the rate of lifetime smoking is 69.5% and girls smoke more than boys. Data obtained in this study show that the rate of smoking among the Turkish students from Turkey has significantly higher than the Cypriot students. In household survey studies conducted in TRNC, They were found that the rate of smoking was 44.7% in 2003 (Çakıcı et al, 2003), 64% in 2008 (Çakıcı et al, 2014) and 62.1% in 2013 (Tutar, 2014). All studies which were conducted in TRNC show that the rate of smoking increases during the transition period between high school and maturity.

1.2.4. Why People Smoke

Studies show that young people smoke for various reasons such as fulfilling their entertainment, social and emotional necessities, getting away from problems, seeking adventure or challenging some negative situations (Hogan, 2000, 27). There are risk factors that encourage young people to smoke, consume alcohol and OPS. These risk factors are family, friends, school, characteristic features of the individual, other risk related behaviors, social and environmental reasons (Ögel, 2002). Mayda et al. (2007) states that the influence of friends on smoking is 54.4%, wannabes 28.0%, curiosity 28.8% and loneliness 20.6%.

1.2.4.1. Biological Reasons

The chemical psychoactive substance that leads to smoking addiction is nicotine and nicotine addictions shows medical similarities to heroine, alcohol and cocaine addiction. It is stated that quitting smoking for a heavy smoker is as hard as giving up heroine for a heroine addict. It is known that by inhaling nicotine reaches the brain within a few seconds, warns several centers and shows its effects. In addition to these, after nicotine delivery is stopped, within 24 hours, symptoms such as an irresistible desire for smoking, uneasiness, anger, anxiety, distractibility, decrease in heart rate and increased appetite are observed. In other words, lack of nicotine causes concrete symptoms known as withdrawal symptom in addicts. Because of these reasons, World Health Organization considers smoking addiction as a disease similar to drug and psychoactive substance use (Dağlı, 1994, 63).

1.2.4.2. Psychological Reasons

Cigarette keeps people from troubles and tension by decreasing the tension experienced in daily life. This situation causes cigarette to become addiction. Behaviors such as taking the cigarette out of the package, lighting up the cigarette, dropping the ash off the cigarette are considered as a way to keep from troubles and tension. It is seen that social and psychological features of youth are important reasons encouraging young people to start smoking and maintaining this habit.

Reasons for smoking initiation are of equivalent value with reasons suggested by various researchers. It is possible to list reasons for smoking as tendency towards estrangement from society, anxiety, stress, influence of friends, curiosity, imitation, wannabe, identification, unlimited autonomy, rebellion against authority, lack of confidence, building relationships with the opposite gender and evading responsibility (Aral and Baran, 1992, 53).

1.2.4.3. Familial Reasons

It is seen that in adolescence period, young people tend to smoke since they want to look like and act as older people. They are unconscious of what this behavior which provides satisfaction in achieving superiority among friends, self-actualization and making himself/herself accepted may cause. Whether parents smoke or not plays a significant role in acquiring this habit. Studies show that young people whose parents smoke show tendency towards smoking more (Mangır et al, 1992, 53). It was found that majority of young people using substance are raised in unhappy families where lack of love, violence and overpermissiveness prevails. Fights between parents, indifference to children, lovelessness and constant domestic tensions injures the mental health of young people driving them into bad environments.

1.2.5. Harms of Cigarette Smoking

It is known that this psychoactive substance is dangerous for health and may cause organic disorder, gastritis, ulcer, lung cancer and heart attack. Furthermore, Abrams (2014) states that cigarette smoking causes many health problems and he lists those problems as such:

- 480.000 deterioration to person, death more than 20 million
- Disorders such as colon, arthritis and blindness
- \$289 billion loss
- 5.6 million premature labor

Donald et al. (1994) states that smoking causes death in the U.S. but people still more than 29% of people continue smoking. In addition to this, as well as physical

and psychological effects, smoking has economic impacts (Sağlık Bakanlığı, 1995, 7). According to studies (Hawkins et al,1992, 64). There needs to be done; In this regard, in order to preserve the mental health of the society, the most important given priority is to fight against substance use within the scope of preventive health services. It is also significant to take care of adolescents who are under the risk of substance addiction by creating opportunities to get to know them and talking to them about the harms of substance use. It can be said that nurses should be trained in terms of substance use and other addictive behaviors.

1.2.6. Treatment of Smoking

When the individual has the need for smoking, he/she should consume nutritious food, water or juice instead of smoking. Furthermore, playing with keys or chain might decrease the desire for smoking. He/she should not stay with smokers for a long time. He/she should jog in the open air, do exercises and pay attention to sports. He/she eat regularly, consume soups, rest for 5-10 minutes after each meal and take a 5-10 minute walk. For cigarette addicts the hardest time of the day is evenings. They should avoid heavy meals, alcohol, coffee, stop watching TV after a while and focus on engrossing things. Continuing this kind of behaviors for a week or two weeks might be the end of smoking habit. The feeling of security after quitting smoking is the best assurance for not starting smoking again.(Aral and Baran, 1992, 53). Ways to quit smoking are listed as follows: Quitting suddenly, quitting slowly, hypnosis, cigarette with low amount of nicotine, gum with nicotine, special cigarette filters and psychological treatment. The most dangerous period after quitting smoking is the first second and third month because 88% of people quitting smoking start smoking within 58 days. One of the most widely used methods for quitting smoking is to take professional help. This kind of help may be provided in various ways. For example;

- Methods based on teaching and conditioning,
- Medicine and science based programs,
- Hypnosis,
- Acupuncture,

- Therapies by restricting environmental stimulants,
- Therapies based on writing imaginary scenarios.

Why do some people restart smoking after quitting? They have valid reasons or excuses for that. These excuses are:

1. "I did not say I would definitely quit smoking. They insisted. Actually, I am strongminded and I can quit smoking any time."
2. "I am not strongminded. I cannot quit smoking. I tried but it did not work."
3. "What difference does it make at all? We are going to die eventually."
4. "Quitting smoking is not good for me. I can quit any time and restart any time."

This kind of excuses are the reasons encouraging people to restart smoking (Barış, 1994). In addition to this, it is known that smoking has physical effects as well as mental and behaviorial effects. When these factors are not investigated sufficiently, it is seen that people restart smoking after nicotine deprivation is over.

It is stated that physical, mental and behaviorial factors are closely related. For instance, craving nicotine (physical addiction) might cause mental problems. Mental problems and depression might increase the desire for nicotine. As in craving tea or coffee while waiting for a bus, behaviorial factors may increase the desire for smoking. It should also be remembered that the number of smokers, encouraging places for smoking, setting other people as examples or adaptation are among reasons for smoking. In addition to this, studies show that behavior consulting ve drug therapy are effective in treatment of smoking cessation (Kaya, 1991, 46).

Medical treatments effective in smoking cessation are nicotine replacement treatment, bupropion and vareniclin. Each treatment has side effects or some cases in which they should not be used (Mangır, 1992, 54). These medicines are prescription drugs, and thus, they should be prescribed by a doctor or used under the supervision of a doctor. Doctors examine their patients attentively, inform patients of the treatment and finally decide on what drug to use (Barut, 1992, 37).

1.3. ALCOHOL

1.3.1. General Information About Alcohol

It is known that alcoholic drinks are psychoactive substances produced from fermented sugary nutrients and these drinks affect the brain and consciousness (Turhan et al, 2011, 34). It is also known that the amount of ethyl alcohol changes and as the amount of alcohol increases, possible damages are more likely to happen. Habitual intoxication which is observed in people who drink heavily is called as alcohol addiction or alcoholism (Yoshimoto, McBride, Lumeng & Li, 1992, 17). Furthermore, it is seen that alcohol addicts are incapable of working gradually, lose his/her job, spend days by drinking alcohol and become a burden on the family and society.

Besides, today there is an increase in the number of people drinking alcohol and becoming addicted. Alcohol is considered as a serious health problem as drugs are also considered as a serious problem. It can also be said that the fact that alcohol is sold freely and celebrities show up with alcoholic drinks on media encourages the use of alcohol. Furthermore, it is not possible to foresee who will become alcohol addict (Hasin, Stinson, Ogburn, & Grant, 2007, 830). For this reason, every drinker is seen as a potential addict. It is also seen that once the habit is gained, it becomes too late because the number of people, who are able to quit drinking as a result of a long and costly treatment period, is significantly less (Turhan et al, 2011, 37).

1.3.2. Use of Alcohol

It is stated that there was a significant increase in the use of alcohol and problems caused by use of alcohol in the last 25-30 years (Turhan et al, 2011, 37). The increase in the use of alcohol in the world in recent years draws a lot of attention. It is stated that the increase is mostly observed especially in developing countries (WHO, 1982, 25). According to data obtained from DSM-IV, the lifelong risk of drinking for women is 10% in America, whereas the lifelong risk of drinking for men is 20%. Also, the lifelong risk of alcohol addiction for women is 3-5%, while it is 10% for men. The prevalence of alcohol addiction and alcohol abuse is reported to be 13,8% (Yavuz et al, 2008, 225). It is also observed that there is an increase in consumption

parallel to production and it was found that 87% of people in America drink alcohol, while 38% of people are addicted to alcohol (Harford, 1992, 32). It was found that alcohol use in Turkey reached 600 million liter in 1992 whereas it was 400 million liter in 1981 (Dağlı, 1994, 63). In addition to this, studies on the use of alcohol in Turkey show that alcohol use increases every day in Turkey.

1.3.3. Alcohol Abuse

It is seen that disorders related to the use of alcohol are categorized into two: alcohol abuse and alcohol addiction (DSM-IV). According to State Institute of Statistics, annual use of alcohol has been increasing significantly (Bayar and Yavuz, 2008, 221). Besides, alcohol use among use is rarely seen in Turkey, but in recent years there has been an increase in the number of women using alcohol in Turkey.

Alcohol abuse is used as a term that is defined by DSM-IV as a stage of alcohol use which has not increased to the level of addiction. Tolerance or deprivation syndrome which is observed when quitting alcohol has not been developed in alcohol abuse as well as using a certain alcohol or consuming too much alcohol (Dağlı, 1994, 63). It is stated that DSM-IV-TR uses the same criteria "psychoactive substance addiction" and "abuse" for every psychoactive substance (Jellinek, 1952, 673). Phases of alcohol addiction. Quarterly journal of studies on alcohol, 13(4), 673-684). It is discussed that the necessity of drinking too much alcohol during the day, consuming too much alcohol regularly on weekends and consuming too much alcohol for a long period of time are indicators of disorders related to alcohol.

Study conducted by Yavuz ve Bayar (2008) shows that young people consuming alcohol come from oppressive families where parents always argue and limitlessness and abuse are constantly experienced. Indifferent, inconsistent and oppressive families increase the rate of drinking (Tol, 1990, 61). Also, another study shows that young people raised in incompatible families tend to drink more alcohol as a reaction against their parents, and even one of the primary reasons for using psychoactive substances is the lack of harmony between parents (Connors et al, 198-247). Besides, according to Didier and Smart; Akfert, Çakıcı and Çakıcı (2009), compared to adolescents with good family relations, adolescents with weak family relations drink more. Çakıcı and Çakıcı (2000) indicate that individuals who are exposed to physical

and psychological abuse and ignorance are more inclined to use psychoactive substances.

In a study by Akfert, Çakıcı and Çakıcı (2009), it was found that the total number of problems faced in families whose children both drink and smoke is significantly high. Also, different studies show that young people having various problems with their families tend to smoke, drink and use psychoactive substances more (Yavuz and Bayar, 2008, 223).

In another study by Combs and Landsverg (1988), it was found that the relationship between parents plays a major role in encouraging adolescents to drink and use psychoactive substances, the communication between young people using psychoactive substances or drinking do not have good relationships with their parents, these adolescents cannot express themselves emotionally, the family has strict rules in terms of doing homeworks, watching TV and so on and these adolescents want to trust their parents and build good relationships with them. It was also found that the parents of adolescents who do not drink and use psychoactive substances reward their children more and help to solve their problems. Youth is a period of time where young people look for identity, show more risky behaviors and the tendency towards smoking and drinking appear in this period more significantly (Turhan et al, 2011, 39). It is known that having a peaceful and happy family atmosphere contributes to the mental development of the adolescents, whereas troubled family atmosphere leads to smoking and drinking. Based on this data, it is significant to note that family plays a major role in smoking and drinking.

1.3.4. History of Alcohol Use

Alcohol is a volatile, pleasure-inducing, depressant, toxic substance that inhibits in the neural system (Kalyoncu ve Mirsal, 2000, 22). Perspectives on alcohol use has changed throughout history (Brown, 2008, 34). It is stated that Bacchus in ancient Rome, Dionysos in Athens were accepted as champagne goddesses, Ancient Egyptians, Jews and Greeks used alcohol in medical interventions and also, these communities confronted dilemmas and rejected alcohol as they realized it caused loss of control. In other words, although alcohol is accepted in every age, excessive use of alcohol is seen as an inappropriate behavior (Köknel, 1998; Brown, 2008). Prehistoric religions used alcohol as a holy token in religious ceremonies. In Judaism

drinking a decent amount of alcohol was seen as a religious activity, whereas Christianity banned drunkenness allowing only drinking. However, although Islam did not intervene in drinking initially, alcohol was banned in Islam afterwards (Brown, 2008, 34). Today, drinking alcohol and liquors is accepted as a part of social interactions in various parts of the world. However, it is seen that alcohol causes social problems due to health problems and the risk of addiction (WHO, 2009, 17).

1.3.5. Prevalence of Alcohol Use in TRNC

In a study conducted by Akfert, Çakıcı ve Çakıcı (2009) in Turkish Republic of Northern Cyprus, the rate of lifelong smoking is 61.5%, whereas the rate of lifelong drinking is 70.8%. It is also seen that these results are consistent with the data obtained from studies in Turkey and Turkish Republic of Northern Cyprus (Yavuz and Bayar, 2008, 225).

In the household survey study conducted in 2003, it was found that drinking alcohol at least once in their lifetime was 82.1%, whereas it was found to be 77.1% in another household survey study conducted in 2008, and 68.5% in household survey study carried out in 2013. In a study by Akfert, Çakıcı and Çakıcı (2009), it was found that 29.7% of students tried smoking at age 18 for the first time, whereas 31.6% of students tried drinking at age 18 for the first time. In the first high school study in TRNC at 1996 which is covered 2515 participant at least once in their life time alcohol use rate was found as %42 (Çakıcı M & Çakıcı E, 2000a). In the four subsequent high school studies alcohol use rates were; %79.7 at 1999, %85.9 at 2004, %75.6 at 2011 and %69.7 at 2015 (Çakıcı M & Çakıcı E, 2000b, Çakıcı et al 2010, Eş 2015, Çakıcı et al., 2015). The results of study conducted with university students showed that the rate of drinking alcohol at least once in their lifetime was 81.0% (Çakıcı et al., 2014). When drinking alcohol in TRNC is compared with alcohol use in Turkey, which share common historical and cultural values (Çakıcı et al., 2014), alcohol use shows differences and the rate of alcohol use in TRNC is higher (Çakıcı et al., 2003).

It is stated that attending a new environment after high school and staying away from the supervision of the society increases the possibility of drinking (Çivi and Şahin, 1991, 49). Furthermore, it is stated in another study that students who try drinking

have bad communication skills in their families although maintaining good communication skills is beneficial for expressing individuals better (Çivi and Şahin, 1991, 49). Also, drinking leads to an increase in selfconfidence, mood and communication skills.

1.3.6. Reasons for Alcohol Use

Many factors play a significant role in starting drinking, having minor problems with alcohol at a young age and having addiction problem at later ages. Related to socio-cultural and psychological problems, drinking is an accepted case in the western society. However, whereas factors influencing drinking lead to temporal problems, they may lead to alcohol addiction in some cases (Schuckit, 2000, 955).

It is not correct to attribute drinking to only one reason. When reasons for drinking are analyzed, it is seen that these reasons are affected by biological, socio-cultural and psychological factors (Yavuz and Bayar, 2008, 223).

1.3.7. Harms of Alcohol Use

Drinking alcohol leads to problems such as hepatitis, liver fattening, cirrhosis and risk of cancer. Drinking has physical risks such as gastritis, esophagitis, pancreatitis, muscle weakness, myolysis, embolism, hypertension, coronary failure, anemia, heart attack and so on. It is also known that drinking causes psychological problems such as depression, sexual problems, mental problems, insomnia, skepticism and addiction (Çivi and Şahin, 1991, 49).

1.3.8. Treatment for Alcohol Use Problems

Alcohol addiction is seen as a personality disorder and in recent years, it is viewed as a disease. While some people drinking alcohol keep it at a social level, others face alcohol abuse and alcohol addiction. Alcohol addiction is regarded as an illness that develops over time and has a destructive effect both on individuals and their family.

Alcohol addiction leads to material and nonmaterial results by causing physical and psychological problems as well as destroying the functionality of the individual. Alcohol addiction is caused by ignoring social activities. Alcohol addict decreases

the amount of time spent with his/her family and friends. It takes time for the alcohol addict to rebuild relationships with family and friends. Usually, close friends and close family members such as parents, children and spouse become aware of the addiction more quickly. For this reason, demand for treatment is usually offered by family members. As problems related to drinking and addiction increase, alcohol addict agrees to receive treatment. Since addiction treatment cannot be achieved without the consent of the addict, the addict should have the will to quit drinking. Alcohol treatment consists of two stages:

- Detoxification
- Psychosocial treatment

Detoxification in Alcohol Addiction: After sustained use of alcohol and quitting or decreasing drinking deprivation (withdrawal symptoms) in relation to withdrawal of alcohol show up. These deprivation symptoms are seen in a wide range from light to heavy. Alcohol withdrawal symptoms may increase to a life threatening point and thus, when the alcohol addict quits drinking, he/she may need to receive medical treatment. Within the first few hours or days after quitting or decreasing drinking, alcohol addict may confront with symptoms as indicated below:

Perspiration

- Increase in pulse rate
- Hand tremor
- Insomnia
- Nausea or vomiting

Psychosocial Treatments in Alcohol Addiction: After detoxification treatment is completed, the psychosocial treatments period covering a long and healthy lifespan begins. The primary objective in this period is to build an alcohol free life and prevent restarting alcohol. Alcohol addiction does not only include alcohol use. It also includes changes in life, social environment and habits of the individual. Therefore, alcohol treatment does not only cover receiving medical treatment but also making significant changes in life style. For this reason, the treatment should be

determined according to the needs of the addict and carried out by a team. As addiction treatment is mainly aimed at changing life style, behaviors and habits, alcohol addict should be willing to quit drinking. Willingness for change makes change possible (Brown, 2008).

Involvement of family members in treatment plays a major role in obtaining positive outcomes. Psychosocial treatments in addiction treatments may last long. As the development of addiction cover a period of time, building an alcohol free life also cover a certain period of time. Alcohol addict may restart drinking in this period. For this reason, it is significant for the addict to realize the importance of the process and requirements necessary for handling this situation. The first step on preventing restarting drinking is to make the addict aware of the high risk situations which may encourage him/her to restart drinking. The individual may face certain situations during the period he/she quits drinking because of various reasons. Separation, health problems, new responsibilities, economic difficulties are challenging and may lead to relapse. Also, social activities such as parties and celebrations pose a risk for restarting drinking. Environment and relatives of the individual may also increase risks. In order to handle this situation, below mentioned methods are suggested:

- Looking for someone who can help,
- Applying methods to solve problems,
- Asking for help from hospitals,
- Joining Alcoholics Anonymous if it is available in the city the alcoholic resides (Turhan et al., 2011, 34).

1.4. PSYCHOACTIVE SUBSTANCE USE AND REASONS

1.4.1. Psychological Reasons

Freud suggests that addiction problems are caused by problems experienced in oral stage and defines oral stage as a period when individual accomplishes psychological development. It is stated that oral personality, excessive dependence on mother and emotional stress develops in this period (Ögel, 1997, 62). Freud also discusses that psychoactive substances such as milk and water are used as a way to satisfy themselves and these substances are replaced by drinking alcohol and smoking in the upcoming stages. Apart from that, Adler supports that incomplete feelings at birth, failures in business life and problems in social life lead to the use of alcohol and drugs. He further states that psychoactive substance such as alcohol and cigarette are used in order to provide self-satisfaction (Allgulander, 1989, 1006).

1.4.2. Genetic Causes

Psychoactive substance use is caused by both environmental and genetic causes. It is observed that the existence of individuals with psychoactive substance use problems creates biological tendency towards psychoactive substance use (Jellinek, 1952, 675). It can also be said that the existence of alcoholism and psychoactive substance use in the family may be effective on the tendency levels of individuals. According to the results of studies, it is seen that genetics plays a major role in the development of alcoholism by 50-70 percent (Ögel, 1997, 63).

1.4.3. Biological Causes

Studies show that some parts and systems of the brain influence addiction. However, Ögel (1997) states that this effect is temporal. Further indicates that body produces endorphin as well as morphine. When body receives drugs, the balance of psychoactive substances change and body needs drugs in order to be able to reshape the balance of these substances (Allgulander, 1989, 1007).


1.4.4. Sociocultural Causes

Psychoactive substances are more accessible gradually which increases the use of psychoactive substance. Ögel (1997) states that the fact that individuals find psychoactive substance use normal increases the prevalence of psychoactive substance use.

1.4.5. Frequency of Psychoactive Substance Use

In 2004, according to a study conducted among 2267 students in 33 high schools in Turkish Republic of Northern Cyprus, the prevalence of life long smoking was 35.2%, the prevalence of lifelong drinking was 85.9% and other psychpactie substance use was 8.0% (Çakıcı et al., 2010, 206). In 2011, according to another study conducted among 2114 students in 34 high schools in Turkish Republic of Northern Cyprus, the prevalence of lifelong smoking was 26.8%, the prevalence of lifelong drinking was 75.6% and prevalence of OPS use was 10.0% (Runeson, 1990, 561). However, in a study conducted among 861 elementary school students aged 13-14, it was found that the prevalence of lifelong smoking was 19.7%, the prevalence of life time drinking was 61,9%, and prevalence of OPS use was 5.8% (Çakıcı et al, 2001, 176). In a study conducted among university students in Turkish Republic of Northern Cyprus, the prevalence of life time smoking was 69.5%, the prevalence of lifelong drinking was 81% and use of OPS was 15.6% (Çakıcı et al., 2014, 159).

There are few studies related to substance use among adults in Turkish Republic of Northern Cyprus. Studies targetting at the society were conducted by using the same technique and survey respectively in 2003, 2008 and 2013. In 2003, according to a study conducted among 825 people in Turkish Republic of Northern Cyprus, it was found that the prevalence of lifelong smoking among 18-65 ages was 44.5%, the prevalence of lifelong drinking was 82.1% and other psychoactive substance use was 5.9% (Çakıcı et al, 2001, 176). In the same way, in another study conducted in 2008 among 804 participants it was found that the prevalence of lifelong smoking was 64%, the prevalence of lifelong drinking was 77.1% and OPS use was 7.7% (Çakıcı et al, 2014, 159). In another study conducted in 2013 among 1040 participants, it was found that the prevalence of lifelong smoking was 62.1%, the prevalence of lifelong



drinking was 82,3% and the other psychoactive substance use was 8.4% (Tutar, 2014). Prevalance studies conducted in Turkish Republic of Northern Cyprus in 2003, 2008 and 2013 show that the most widely used substance is drugs. Also, studies conducted in the time period show that use of bonsai, ecstasy, codein syrup and calmatives are quite common. Especially the increase in the use of bonsai in Turkish Republic of Northern Cyprus in recent years is dramatic (Tutar, 2014, 14). It's seen that the fact that Turkish Republic of Northern Cyprus is a small country makes the access of psychoactive substances into the country possible and increase the use of these substances (Çakıcı et al, 2001, 177). Use of drugs has been increasing day by day and the drug age has been decreasing significantly (Tutar, 2014,14). M. Çakıcı and E. Çakıcı (2000a) states that drug use has a negative impact on young people. Ögel (1997) indicates that America is the country where the highest amount of drugs are produced and mariuana is the substance that is used most in these countries. It can be said that psychoactive substance use has a negative impact on the health of society. Furthermore, Köknel (1998) states that the use of alcohol, smoking and drugs has been increasng in developing countries in Middle East. The psychoactive substance most widely used in Turkey is cigarette (Tutar, 2014, 15). Besides, it is seen that psychoactive substance use in many countries such as Turkey where level of education and income is low has been increasing gradually (Ögel, 1997, 65). Ögel and Başterzi (2010) states that access to psychoactive substances and cheap prices make these substances attractive. Thus, it can be said that these reasons lead to increase in the use of psychoactive substances.

2. METHOD

2.1. Sample

The study took place among Turkish speaking individuals at age 18-65 in Turkish Republic of Northern Cyprus. The study was conducted among 994 participants in Turkish Republic of Northern Cyprus and based on quota multiple-stage randomized sample. Individuals participating in the study are chosen based on gender (male, female), age (18-19, 20-29, 30-39, 40-49, 50-65), settlement (village, city) and features of the region where individuals live. The choice of sample is based on the statistics obtained from population census on 4 December, 2011 (Population Census, 2011). In the light of the results obtained from the last population census, population characteristics in five regions including Nicosia, Famagusta, Kyrenia, Guzelyurt and Iskele were taken into consideration. These 5 regions were divided into neighborhoods in cities and villages in rural regions. This kind of randomized study included 16 neighborhoods, 17 villages and sub-districts (Lefke, Güzelyurt, Mehmetçik, İskele, Geçitkale).

2.2. Survey

2.2.1. The Model European Questionnaire

Survey was prepared by considering "The Model European Questionnaire." (EMCDDA, 1995). European Monitoring Centre for Drugs and Drug Addiction EMCDDA (1995) was taken from the study conducted by Çakıcı et al. (2003, 2008 ve 2013). Model European Questionnaire has never been applied before the study done by Çakıcı et al (2003) to the Turkish Cypriot population. In the Çakıcı et al. (2003) study, original survey questionnaire is translated by one academician from Education Faculty, two academician from Art and Sciences Faculty in total of three academicians of Near East University. Then it is translated from Turkish to English by an academician who is attendant in English Language Department, after that one of the academician who is from English Language Department is decided that every question is qualified. Survey form includes informed consent and sociodemographic forms. The Model European Questionnaire includes two parts which

help to design of substance use prevalence. First part of the survey covers sociodemographic questions like age, sex, social support, income level, work life, education level. Second part of survey covers questions which help to find out frequency of substance use, reasons for substance use. By use of this survey it is aimed to compare with other European countries.

2.3. Procedure

Study was conducted in Turkish Republic of Northern Cyprus in May-June 2015. Starting points were determined randomly as streets, villages and main spots in villages (tea houses or mosque) and covered north, south, east and west.

Pollsters started from the right side of the road and the minimum numbers by creating a square shape. One in every three houses was included in the study and the route of the study was determined by turning right in every street and creating a square shape. When one square was completed, another square shape was restarted from the street situated below the starting point. This way it was ensured that mistakes could be prevented by making a mutual point in the choice of the houses selected by the pollsters. One in every three houses was included in the study and males and females along with their ages were taken into consideration. Only one person in the house visited was included in the study. The study was conducted in a way that one male in the first house and one female in the second house participated in the survey. In terms of age quota, if there were more than one person in the house, the one whose birthday was approaching was selected. Survey is completed by participant and it is collected in a closed box. 47 pollsters took part in the study and pollsters received training before attending the study. Every pollster conducted a poll with 21 people at most. This way it was aimed to decrease the possibility of error margin.

2.4. Statistical Method

Descriptive statistics data was used in the study. Comparison of sociodemographic characteristics and OPS use of the male and female differences and also tobacco and alcohol user and non-user participants differences chi-squared statistic method was used. For investigating the relationship between risk factors and cigarette and alcohol use multivariate logistic regression analysis was used.

3. RESULT

As a result of the survey 994 participants were analysed. 490 participants were female and 504 participant were male. Age distribution of the participants were 30.9% 18-29 age group, 22.8% 30-39 age group, 19.3% 40-49 age group, 14% 50-59 age group and 13% 60 and over age group.

Table 1. Comparison of male and female distribution of participants in terms of smoking cigarette, pipe or cigar

	Female		Male		Total	
	N	%	N	%	N	%
Smokers	163	33.5	246	50.1	409	41.8
Nonsmokers	324	66.5	245	49.9	569	58.2
Total	487	100.0	491	100.0	978	100.0

$X^2=27.796$, $df=1$, $p=0.000$, DA (Do not answer)=16 (%1.6)

There is statistically significant difference between males and females for smoking cigarette, pipe or cigar according to chi-square statistical method. Male participants smoke more than female participants ($X^2=27.796$, $df=1$, $p=0.000$).

Table 2. Comparison of male and female distribution of the time of serving the highest amount of alcohol in participant houses

	Female		Male		Total	
	N	%	N	%	N	%
Never	194	41.4	179	37.5	373	39.4
When guests come over	108	23.0	119	24.9	227	24.0
Any time, without appetizer, as a relaxing drink	9	1.9	23	4.8	32	3.4
Any time with appetizer	15	3.2	30	6.3	45	4.8
At lunch	0	0.0	0	0.0	0	0.0
At dinner	9	1.9	18	3.8	27	2.9
Only on Sundays	9	1.9	7	1.5	16	1.7
While celebrating something	109	23.2	77	16.1	186	19.7
Other	16	3.4	24	5.0	40	4.2
Total	469	100.0	477	100.0	946	100.0

$X^2 = 22.551$, $df = 7$, $p = 0.002$, $DA = 48$ (%4.8)

There is statistically significant difference between males and females for the time of serving the highest amount of alcohol in participant houses according to chi-square statistical method. Females drink when guests come or they celebrate something, but males drink more alcohol when guests come, dinners, for relaxing time and they celebrate something

Table 3. Comparison of male and female age distribution of participants to start smoking

Age	Female		Male		Total	
	N	%	N	%	N	%
11 and below	6	2.8	18	5.2	24	4.3
12	3	1.4	18	5.2	21	3.8
13	5	2.3	11	3.2	16	2.9
14	8	3.7	23	6.7	31	5.6
15	12	5.6	31	9.0	43	7.7
16	16	7.5	33	9.6	49	8.8
17	21	9.8	31	9.0	52	9.3
18 and above	143	66.8	178	51.9	321	57.6
Total	214	100.0	343	100.0	557	100.0

$X^2=17.307$, $df=7$, $p=0.016$, $DA=437$ (%44.0)

There is statistically significant difference between males and females for age to start according to chi-square statistical method. Males have started smoking more below the age of 11 ($X^2=17.307$, $df=7$, $p=0.016$).

Table 4. Comparison of male and female age distribution of participants in terms of drinking constantly more than their peers

Age	Female		Male		Total	
	N	%	N	%	N	%
11 and below	1	2.6	3	2.0	4	2.2
12	0	0.0	1	0.7	1	0.5
13	0	0.0	1	0.7	1	0.5
14	2	5.3	1	0.7	3	1.6
15	1	2.6	6	4.1	7	3.8
16	3	7.9	10	6.8	13	7.0
17	2	5.3	10	6.8	12	6.5
18 and above	29	76.3	115	78.2	144	77.8
Total	38	100.0	147	100.0	185	100.0

$X^2=4,820$, $df=7$, $p=0,682$, $DA=395$ (%39.7)

There is no statistically significant difference between males and females for age to start drinking constantly more than their peers according to chi-square statistical method. Mostly both females and males are have started drinking constantly after 18 and above ($X^2=4,820$, $df=7$, $p=0,682$).

Table 5. Comparison of male and female age distribution of participants in terms of beginning to drink

Age	Female		Male		Total	
	N	%	N	%	N	%
11 and below	3	1.4	6	1.6	9	1.5
12	2	0.9	10	2.6	12	2.0
13	0	0.0	8	2.1	8	1.3
14	3	1.4	14	3.7	17	2.8
15	9	4.1	51	13.4	60	10.0
16	9	4.1	41	10.7	50	8.3
17	25	11.5	36	9.4	61	10.2
18 and above	166	76.5	216	56.5	382	63.8
Total	217	100.0	382	100.0	599	100.0

$X^2 = 37.234$, $df=7$, $p=0.000$, $DA=809$ (81.4)

There is statistically significant difference between males and females for age of beginning to drink according to chi-square statistical method. Comparing males and females males have started drinking below the age of 18 more than females.

($X^2 = 37.234$, $df=7$, $p=0.000$).

Table 6. Comparison of male and female distribution of life long use of smoking

	Female		Male		Total	
	N	%	N	%	N	%
0	435	48.5	128	26.1	363	37.3
1-2	39	8.1	24	4.9	63	6.5
3-5	27	5.6	13	2.7	40	4.1
6-9	6	1.2	5	1.0	11	1.1
10-19	8	1.7	4	0.8	12	1.2
20-39	8	1.7	9	1.8	17	1.7
40 and more	161	33.3	307	62.7	468	48.0
Total	484	100.0	490	100.0	974	100.0

$X^2 = 87.5$, $df=7$, $p=0.000$, $DA = 437$ (%44.0)

There is statistically significant difference between males and females comparison of lifetime use of smoking according to chi-square statistical method. The rate of those who smoked 40 times or more is 33% among females and 62% among males

($X^2 = 87.5$, $df=7$, $p=0.000$).

Table 7. Comparison of male and female distribution of smoking in the past 12 months

	Female		Male		Total	
	N	%	N	%	N	%
0	312	64.5	219	44.5	531	54.4
1-2	15	3.1	9	1.8	24	2.5
3-5	6	2.2	12	2.4	18	1.8
6-9	8	1.7	4	0.8	12	1.2
10-19	5	1.1	4	0.8	9	0.9
20-39	11	2.3	11	2.2	22	2.3
40 and more	127	26.2	233	47.4	360	36.9
Total	484	100.0	492	100.0	976	100.0

$X^2 = 53.063$, $df=7$, $p=0.000$, $DA = 18$ (%1.8)

There is statistically significant difference between males and females comparison of smoking in the past 12 months according to chi-square statistical method. Males participants stated that they smoked more than females ($X^2=53.063$, $df=7$, $p=0.000$).

Table 8. Comparison of male and female distribution smoking in the past 30 days

	Female		Male		Total	
	N	%	N	%	N	%
Hiç içmeyenler	330	68.9	230	46.9	560	57.8
Less than one cigarette a week	14	2.9	17	3.5	31	3.2
Less than a cigarette a day	4	0.8	6	1.2	10	1.0
1-5 cigarettes a day	34	7.1	21	4.3	55	5.7
6-10 cigarettes a day	25	5.2	18	3.7	43	4.4
11-20 cigarettes a day	32	6.7	67	13.7	99	10.2
20-30 cigarettes a day	17	3.5	54	11.0	71	7.3
31-40 cigarettes a day	3	0.6	14	2.9	17	1.8
More than 40 cigarettes a day	20	4.2	63	12.9	83	8.6
Total	479	100.0	490	100.0	969	100.0

$X^2= 83.696$, $df=8$, $p=0.000$, $DA=25$ (%2.5)

There is statistically significant difference between males and females comparison of smoking in the past 30 months according to chi-square statistical method. Males have smoked more than females for more than 40 cigarettes a day in last 30 days ($X^2= 83.696$, $df=8$, $p=0.000$).

Table 9. Comparison of participants in terms of whether they faced difficulty in quitting smoking or not

	Female		Male		Total	
	N	%	N	%	N	%
0	258	53.9	127	25.6	385	39.4
1-2	57	11.7	27	5.4	83	8.5
3-5	35	7.3	35	7.0	70	7.2
6-9	32	6.7	25	5.0	57	5.8
10-19	37	7.7	62	12.5	99	10.1
20-39	20	4.2	40	8.0	60	6.1
40 and more	41	8.6	181	36.4	222	22.7
Total	479	100.0	497	100.0	976	100.0

$X^2=11.109$, $df=3$, $p=0.011$, $DA=323$ (%32.5)

There is statistically significant difference between males and females comparison of participants in terms of whether they faced difficulty in quitting smoking or not according to chi-square statistical method. Males have faced more difficulties in quitting smoking than females ($X^2=11.109$, $df=3$, $p=0.01$).

Table 10. Comparison of participants in terms of consuming alcoholic drinks in the past 12 months

	Female		Male		Total	
	N	%	N	%	N	%
0	258	53.9	127	25.6	385	39.4
1-2	57	11.7	27	5.4	83	8.5
3-5	35	7.3	35	7.0	70	7.2
6-9	32	6.7	25	5.0	57	5.8
10-19	37	7.7	62	12.5	99	10.1
20-39	20	4.2	40	8.0	60	6.1
40 and more	41	8.6	181	36.4	222	22.7
Total	479	100.0	497	100.0	976	100.0

$X^2=156.556$, $df= 6$, $p= 0.000$, $DA=18$ (%1.8)

There is statistically significant difference between males and females comparison of participants distribution of participants in terms of consuming alcoholic drinks in the past 12 months according to chi-square statistical method. Male participants drank more alcohol than female participants in the past 12 months ($X^2=156.556$, $df=6$, $p=0.000$).

Table 11. Comparison of participants in terms of consuming alcoholic drinks in the past 30days

	Female		Male		Total	
	N	%	N	%	N	%
0	329	58.4	184	37.7	513	52.7
1-2	72	15.0	73	14.8	145	14.9
3-5	32	6.7	53	10.8	85	8.7
6-9	15	3.1	36	7.3	51	5.2
10-19	13	2.7	50	10.1	63	6.5
20-39	10	2.1	27	5.5	37	3.8
40 and more	10	2.1	70	14.2	80	8.2
Total	481	100.0	493	100.0	974	100.0

$X^2=129.239$, $df=6$, $p=0.000$, $DA=20$ (%2.0)

There is statistically significant difference between males and females comparison of participants in terms of consuming alcoholic drinks in the past 30days according to chi-square statistical method. Female participants consumed less alcohol than male participants in last 30 days ($X^2=129.239$, $df=6$, $p=0.000$).

Table 12. Comparison of participants in terms of consuming alcohol in the past 30 days

	Female		Male		Total	
	N	%	N	%	N	%
Never	336	71.2	196	39.8	532	55.2
More than twice a week	25	5.3	77	15.7	102	10.6
Once two weeks	70	14.8	85	17.3	155	16.1
Once a week	25	5.3	65	13.2	90	9.3
Once a day	1	0.2	32	6.5	33	3.4
Twice a week	15	3.2	37	7.5	52	5.4
Total	472	100.0	492	100.0	964	100.0

$X^2=120.647$, $df=5$, $p=0.000$, $DA=30$ (%3.0)

There is statistically significant difference between males and females comparison of participants in terms of consuming alcoholic drinks in the past 30 days according to chi-square statistical method. Male participants have drunk more alcohol than females in last 30 days. Majority of female participants did not drink alcohol in the past 30 days ($X^2=120.647$, $df=5$, $p=0.000$).

Table 13. Comparison of the amount of alcohol consumed by participants in one go (one drink, a bottle or a glass of beer, a glass of champagne, a glass of raki or other alcoholic drinks)

	Female		Male		Total	
	N	%	N	%	N	%
Never	227	47.9	99	20.0	326	33.7
1-2 glass	201	42.4	199	40.3	400	41.3
3-4 glasses	36	7.6	126	25.5	162	16.7
5 or more drinks	10	2.1	70	14.2	80	8.3
Total	474	100.0	494	100.0	968	100.0

$X^2=144.916$, $df=3$, $p=0.000$, $DA=26$ (%2.6)

There is statistically significant difference between males and females comparison of the amount of alcohol consumed by participants in one go (one drink, a bottle or a glass of beer, a glass of champagne, a glass of raki or other alcoholic drinks) according to chi-square statistical method. 1-2 glass of alcohol drink in one go are similar in both males and females But 3 glass of more drink in one go are more in males compare with females ($X^2=144.916$, $df=3$, $p=0.000$).

Table 14. Comparison of last places where participants are being drunk

	Male		Female		Total	
	N	%	N	%	N	%
Non-drinkers	164	36.9	70	14.4	234	25.1
At home	117	26.3	151	31.1	268	28.8
At someone else's house	16	3.6	30	6.2	46	4.9
Outside, At a park or in the street	21	4.7	56	11.5	77	8.3
At a bar or cafe	42	9.4	55	11.3	97	10.4
At a disco	7	1.6	22	4.5	29	3.1
At a diner	67	15.1	79	16.3	146	15.7
Other	11	2.5	23	4.7	34	3.7
Total	445	100.0	486	100.0	931	100.0

$X^2=75.307$, $df=7$, $p=0.000$, $DA=63$ (%6.3)

There is statistically significant difference between males and females comparison of last places where participants are being drunk according to chi-square statistical method. Comparing females and males, females have preferred to use alcohol at outside Males have preferred to use at home and dinners ($X^2=75.307$, $df=7$, $p=0.000$).

Table 15. Comparison of participants in terms of how many times they got drunk in their lives

	Male		Female		Total	
	N	%	N	%	N	%
0	186	37.5	337	71.1	523	53.9
1-2	104	21.0	70	14.8	174	17.9
3-5	75	15.1	40	8.4	115	11.9
6-9	34	6.9	6	1.3	40	4.5
10-19	28	5.6	9	1.9	37	3.8
20-39	9	1.8	6	1.3	15	1.5
40 or more	60	12.1	6	1.3	86	6.8
Total	496	100.0	474	100.0	970	100.0

$X^2=134.601$, $df=6$, $p=0.000$, $DA=24$ (%2.4)

There is statistically significant difference between males and females comparison of being drunk in their life according to chi-square statistical method. Males have become more drunk than female ($X^2=134.601$, $df=6$, $p=0.000$).

Table 16. Comparison of participants being drunk in the past 12 months

	Male		Female		Total	
	N	%	N	%	N	%
0	396	65.3	316	85.5	712	75.1
1-2	46	15.1	73	9.9	119	12.1
3-5	8	6.8	33	1.7	41	4.3
6-9	5	2.3	13	1.1	18	1.9
10-19	1	3.3	16	0.2	17	1.8
20-39	5	2.3	11	1.1	16	1.7
40 or more	42	4.5	22	0.4	24	0.5
Toplam	463	100.0	484	100.0	947	100.0

$X^2=65.633$, $df=6$, $p=0.000$, $DA=47$ (%4.7)

There is statistically significant difference between males and females comparison of participants being drunk in the past 12 months according to chi-square statistical method. Males have become more drunk than female in last 12 months ($X^2=65.633$, $df=6$, $p=0.000$).

Table 17. Comparison of participants being drunk in the past 30 days

	Female		Male		Total	
	N	%	N	%	N	%
0	438	94.8	377	78.4	815	86.4
1-2	14	3.0	60	12.5	74	7.8
3-5	5	1.1	18	3.7	23	2.4
6-9	3	0.6	12	2.5	15	1.6
10-19	2	0.4	16	1.2	8	0.8
20-39	0	0.0	3	0.6	3	0.3
40 and more	0	0.0	5	1.0	5	0.5
Total	462	100.0	481	100.0	943	100.0

$X^2=55.548$, $df=6$, $p=0.000$, $DA=51$ (%5.1)

There is statistically significant difference between males and females comparison of participants being drunk in the past 30 days according to chi-square statistical method. have become more drunk than female in last 12 months ($X^2=55.548$, $df=6$, $p=0.000$).

Table 18. Comparison of reasons why participants got drunk

	Female		Male		Total	
	N	%	N	%	N	%
Entertaining	208	75.6	295	72.8	503	74.0
Sleeping	2	0.7	11	2.7	13	1.9
Trying	16	5.8	7	1.7	23	3.4
Being angry	4	1.5	6	1.5	10	1.5
Stress	7	2.5	15	3.7	22	3.2
Relaxing	9	3.3	29	7.2	38	5.6
Staying away from problems	1	0.4	2	0.5	3	0.4
Friends drinking alcohol	19	6.9	28	6.9	47	6.9
Feeling good	9	3.3	12	3.0	21	3.1
Total	275	100.0	405	100.0	680	100.0

$X^2=16.885$, $df=8$, $p=0.031$, $DA=314$ (%31.6)

There is statistically significant difference between males and females comparison of reasons why participants got drunk according to chi-square statistical method. Both female and male participants have drunk alcohol mostly for entertaining. Female participants have drunk alcohol for trying and feeling good male participants consumed alcohol for relaxing and stress ($X^2=16.885$, $df=8$, $p=0.031$).

Table 19. Comparison of gender distribution of smokers and non-smokers

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Male	163	39.9	324	56.9	487	49.8
Female	246	60.1	245	43.1	491	50.2

$X^2=27.796$, $df=1$, $p=0.000$, Don't answer (DA)=16(%1.6)

There is statistically significant difference between smokers and non-smokers of comparison of gender distribution according to chi-square statistical method. Among smokers, males are found to be more than females ($X^2=27.796$, $df=1$, $p=0.000$).

Table 20. Comparison to place of birth distribution of smokers and non-smokers

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Cyprus	241	58.6	357	62.9	598	61.1
Turkey	159	38.7	197	34.7	356	36.4
England	5	1.2	3	0.5	8	0.8
Other	6	1.5	11	1.9	17	1.7

$X^2=3.439$, $df=3$, $p=0.329$, DA=15(%1.5)

There is no statistically significant difference between smokers and non-smokers of comparison to place of birth distribution according to chi-square statistical method. Both Cypriots and Turkish citizens ratio of smoke have seen similarly ($X^2=3.439$, $df=3$, $p=0.329$).

Table 21. Comparison to age distribution of smokers and non-smokers

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
1.18-25	129	31.4	171	30.4	300	30.8
2.26-35	100	24.3	123	21.8	223	22.9
3.36-45	84	20.4	104	18.5	188	19.3
4.46-55	57	13.9	80	14.2	137	14.1
5 56-and more	41	10.0	85	15.1	126	12.9
TOTAL	411	100.0	563	100.0	974	100.0

$X^2=6.032$, $df=4$, $p=0.197$, $DA=20(\%2.0)$

There is no statistically significant difference between smokers and non-smokers of age distribution according to chi-square statistical method. Smokers aged between 18 and 25 smoke more. In other words, it is seen that smoking at a young age is more common ($X^2=6.032$, $df=4$, $p=0.197$).

Table 22. Comparison of smokers and non-smokers in terms of who they live with

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Alone	47	11.7	48	8.6	95	9.9
With partner or spouse	192	47.8	333	59.8	525	54.7
With children	29	7.2	25	4.5	54	5.6
With parents	73	18.2	99	17.8	172	17.9
Mother or father	22	5.5	17	3.1	39	4.1
Step mother/step father	1	0.2	0	0.0	1	0.1
Relatives	8	2	10	1.8	18	1.9
Friends	19	4.7	16	2.9	35	3.6
Other	11	2.7	9	1.6	20	2.1
Total	402	100.0	557	100.0	959	100.0

$X^2=19.894$, $df=8$, $p=0.011$, $DA=35(\%3.5)$

There is statistically significant difference between smokers and non-smokers in terms of who they live with according to chi-square statistical method. Smokers live alone at a higher rate than non-smokers, non-smokers live with partner or spouse ($X^2=19.894$, $df=8$, $p=0.011$).

Table 23. Comparison of education level distribution of smokers and non-smokers

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Non-schoolers	11	2.7	17	3.0	28	2.9
Primary school	84	20.7	124	21.9	208	21.4
Elementary school	60	14.8	57	10.1	117	12.1
High school	133	32.8	184	32.6	317	32.7
University and more	117	28.9	183	32.4	300	30.9
Total	405	100.0	565	100.0	970	100.0

$X^2=5.539$, $df=4$, $p=0.236$, $DA=24(\%2.4)$

There is no statistically significant difference between smokers and non-smokers of education level according to chi-square statistical method. It is seen that smoking in high school and university level is more common ($X^2=5.539$, $df=4$, $p=0.236$).

Table 24. Comparison of education level distribution of fathers of smokers and non-smokers

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Non-schoolers	62	15.2	100	17.8	162	16.7
Primary school	190	46.6	268	47.8	458	47.3
Elementary school	59	14.5	77	13.7	136	14.0
High school	62	15.2	78	13.9	140	14.4
University and more	35	8.6	38	6.8	73	7.5
Total	408	100.0	561	100.0	969	100.0

$X^2=2.434$, $df=4$, $p=0.656$, $DA=25(\%2.5)$

There is no statistically significant difference between comparison of education level distribution of fathers of smokers and non-smokers according to chi-square statistical method. It is seen that the education level of fathers of smokers is primary school. According to this table, individuals whose fathers are university graduates smoke more ($X^2=2.434$, $df=4$, $p=0.656$).

Table 25. Comparison of education level distribution of mothers of smokers and non-smokers

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Non-schoolers	90	22.1	121	21.5	211	21.7
Primary school	173	42.4	266	47.2	439	45.2
Elementary school	49	12.0	58	10.3	107	11.0
High school	75	18.4	90	16.0	165	17.0
University and more	21	5.1	28	5.0	49	5.0
Total	408	100.0	563	100.0	971	100.0

$X^2=2.703$, $df=4$, $p=0.609$, $DA=23(\%2.3)$

There is no statistically significant difference between comparison of education level distribution of mothers of smokers and non-smokers according to chi-square statistical method. According to this table, individuals whose mothers are primary school graduates smoke more ($X^2=2.703$, $df=4$, $p=0.609$).

Table 26. Comparison to birth place distribution to mothers of smokers and nonsmokers

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Cyprus	211	51.6	322	56.6	533	54.5
Turkey	187	45.7	232	40.8	419	42.8
England	3	0.7	1	0.2	4	0.4
Other	8	0.8	14	2.5	22	2.2
Total	409	100.0	569	100.0	978	100.0

$X^2=4.531$, $df=3$, $p=0.210$, $DA=16(\%1.6)$

There is no statistically significant difference between comparison to birth place distribution of mothers of smokers and non-smokers according to chi-square statistical method. Mothers of smokers are mostly born in Cyprus and Turkey ($X^2=4.531$, $df=3$, $p=0.210$).

Table 27. Comparison to birth place distribution of fathers of smokers and non-smokers

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Cyprus	199	48.7	318	55.9	517	52.9
Turkey	200	48.9	232	40.8	432	44.2
England	1	0.1	3	0.5	4	0.4
Other	9	2.2	16	2.8	25	2.6
Total	409	100.0	569	100.0	978	100.0

$X^2=6.725$, $df=3$, $p=0.081$, $DA=16(\%1.6)$

There is no statistically significant difference between comparison to birth place distribution of fathers of smokers and non-smokers according to chi-square statistical method. According to this table, fathers of heavy smokers are mostly born in Cyprus and Turkey ($X^2=6.725$, $df=3$, $p=0.081$).

Table 28. Comparison of distribution to places where smokers and non-smokers live

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Village	151	36.8	266	46.9	417	42.7
City	241	58.8	291	51.3	532	54.5
Suburs	18	4.4	10	1.8	28	2.9
Total	410	100.0	567	100.0	977	100.0

$X^2=13.827$, $df=2$, $p=0.001$, $DA=17(\%1.7)$

There is statistically significant difference between comparison of distribution to places where smokers and non-smokers live according to chi-square statistical method. People living in cities smoke more compared to those living in villages and suburbs ($X^2=13.827$, $df=2$, $p=0.001$).

Table 29. Comparison of work status distribution of smokers and non-smokers

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Yes	271	66.9	295	52.8	566	58.7
Non	134	33.1	264	47.2	398	41.3
Total	405	100.0	559	100.0	964	100.0

$$X^2=19.373, df=1, p=0.000, DA=30(\%3.0)$$

There is statistically significant difference between comparison of work status distribution of smokers and non-smokers according to chi-square statistical method. Compared to unemployed participants, employed participants smoke more ($X^2=19.373, df=1, p=0.000$).

Table 30. Comparison of social support distribution of smokers and non-smokers

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Yes	149	36.8	242	43.4	391	40.6
No	253	62.5	315	56.5	568	59.0
Total	405	100.0	558	100.0	963	100.0

$$X^2=5.724, df=2, p=0.057, DA=35(\%3.7)$$

There is no statistically significant difference between comparison of social support distribution of smokers and non-smokers according to chi-square statistical method. It is seen that those who do not receive social support smoke more ($X^2=5.724, df=2, p=0.057$).

Table 31. Comparison of income status distribution of smokers and non-smokers

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Very good	10	2.4	12	2.1	22	2.3
Good	126	30.7	194	34.2	320	32.8
Average	234	57.1	328	57.8	562	57.5
Bad	30	7.3	26	4.6	56	5.7
Very bad	10	2.4	7	1.2	17	1.7
Total	410	100.0	567	100.0	977	100.0

$$X^2=6.098, df=4, p=0.192, DA=17(\%1.7)$$

There is no statistically significant difference between comparison income status distribution of smokers and non-smokers according to chi-square statistical method. It can be said that income status does not influence smoking ($X^2=6.098$, $df=4$, $p=0.192$).

Table 32. Comparison of smokers and non-smokers in terms of where they receive most information about smoking

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Media	284	71.4	432	77.7	716	75.1
Book,newspaper,booklet	17	4.3	22	4.0	39	4.1
Family and friends	7	1.8	14	2.5	21	2.2
Friends	45	11.3	24	4.3	69	7.2
Teachers	0	0.0	5	0.9	5	0.5
Doctor, nurse or health officers	6	1.5	4	0.7	10	1.0
Youth centers	1	0.3	0	0.0	1	0.1
Internet	18	4.5	37	6.7	55	5.8
Other	20	5.0	18	3.2	38	4.0
Total	398	100.0	556	100.0	954	100.0

$X^2=27.617$, $df=8$, $p=0.001$, $DA=40(\%4.0)$

There is statistically significant difference between comparison of smokers and non-smokers in terms of where they receive most information about smoking according to chi-square statistical method. It is seen that non-smokers receive information about smoking mostly from media, smokers receive information from friends ($X^2=27.617$, $df=8$, $p=0.001$).

Table 33. Comparison of smokers and non-smokers in terms of the importance of religion in their lives

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Very important	187	47.7	267	49.7	454	48.9
Relatively important	155	39.5	218	40.6	373	40.2
Unimportant	50	12.8	52	9.7	102	11.0
Total	392	100.0	537	100.0	929	100.0

$X^2=2.199$, $df=2$, $p=0.333$, $DA=65(\%6.5)$

There is no statistically significant difference between comparison of smokers and non-smokers in terms of the importance of religion in their lives according to chi-square statistical method. It is seen that religion does not influence smoking significantly ($X^2=2.199$, $df=2$, $p=0.333$).

Table 34. Comparison of gender distribution of drinkers and non-drinkers

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Male	424	47.7	70	87.5	494	51.0
Female	464	52.3	10	12.5	474	49.0
Total	888	100.0	80	100.0	968	100.0

$X^2=46.408$, $df=1$, $p=0.000$, $DA=26(\%2.6)$

There is statistically significant difference between comparison of gender distribution of drinkers and non-drinkers according to chi-square statistical method. Statistically significant difference has been found. Male participants drink more than female participants ($X^2=46.408$, $df=1$, $p=0.000$).

Table 35. Comparison of birthday distribution of drinkers and non-drinkers

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
18-25	265	30.0	37	46.3	302	31.3
26-35	202	22.9	20	25.0	222	23.0
36-45	173	19.6	12	15.0	185	19.2
46-55	123	13.9	8	10.0	131	13.6
56 and more	121	13.7	3	3.8	124	12.9
Total	884	100.0	80	89.0	964	100.0

$X^2=13.606$, $df=4$, $p=0.009$, $DA=30(\%3.0)$

There is statistically significant difference between comparison of birthday distribution of drinkers and non-drinkers according to chi-square statistical method. There is a statistical difference between both groups. It is seen that participants aged between 18 and 25 drink more. It is also observed that drinking is more widely seen among young participants than older participants ($X^2=13.606$, $df=4$, $p=0.009$).

Table 36. Comparison of distribution to drinkers and non-drinkers in terms of who they live with

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Alone	88	10.1	10	12.7	98	10.3
With partner or spouse	485	55.0	28	35.4	513	54.0
With children	52	6.0	5	6.3	57	6.0
With parents	148	17.0	22	27.8	170	17.9
With mother or father	34	3.9	5	6.3	39	4.1
With step mother/step father	1	0.1	0	0.0	1	0.1
With relatives	14	1.6	4	5.1	18	1.9
With friends	29	3.3	4	5.1	33	3.5
Other	20	2.3	1	1.3	21	2.2
Total	871	100.0	79	100.0	950	100.0

$X^2=17.408$, $df=8$, $p=0.026$, $DA=44(\%4.4)$

There is statistically significant difference between comparison of distribution of drinkers and non-drinkers in terms of who they live with according to chi-square statistical method. Drinkers live with their parents, non-drinkers live with partner or spouse ($X^2=17.408$, $df=8$, $p=0.026$).

Table 37. Comparison of education level distribution of drinkers and non-drinkers

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Non-schoolers	25	2.8	1	1.3	26	2.7
Primary school	190	21.6	14	17.5	204	21.2
Elementary school	105	11.9	11	13.8	116	12.1
High school	291	33.0	26	32.5	317	33.0
University and more	270	30.6	28	35.0	298	31.0
Total	881	100.0	80	100	961	100.0

$X^2=1.913$, $df=4$, $p=0.752$, $DA=33(\%3.3)$

There is no statistically significant difference between comparison of education level distribution of drinkers and non-drinkers according to chi-square statistical method. It is seen that participants graduating from high school and university drink more than other participants ($X^2=1.913$, $df=4$, $p=0.752$).

Table 38. Comparison of education level distribution of fathers to drinkers and non-drinkers

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Non-schoolers	145	16.5	15	18.8	160	16.7
Primary school	424	48.2	24	30.0	448	46.7
Elementary school	119	13.5	16	20.0	135	14.1
High school	127	14.4	15	18.8	142	14.8
University and more	64	7.3	10	12.5	74	7.7
Total	879	100.0	80	100.0	959	100.0

$X^2=11.123$, $df=4$, $p=0.025$, $DA=35(\%3.5)$

There is statistically significant difference between comparison of education level distribution of fathers of drinkers and non-drinkers according to chi-square statistical method. Participants whose fathers are primary school graduate smoke more. As the education level of fathers decrease, drinking increases. ($X^2=11.123$, $df=4$, $p=0.025$).

Table 39. Comparison of education level distribution of mothers of drinkers and non-drinkers

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Non-schoolers	191	21.7	17	21.5	208	21.6
Primary school	404	45.8	25	31.6	429	44.6
Elementary school	97	11.0	11	13.9	108	11.2
High school	146	16.6	20	25.3	166	17.3
University and more	44	5.0	6	7.6	50	5.2
Total	882	100.0	79	100.0	961	100.0

$X^2=7.979$, $df=3$, $p=0.092$, $DA=33(\%3.3)$

There is no statistically significant difference between comparison of education level distribution of mothers of drinkers and non-drinkers according to chi-square statistical method. According to this table, individuals whose mothers are primary school graduates smoke more ($X^2=7.979$, $df=3$, $p=0.092$).

Table 40. Comparison of birth place distribution of drinkers and non-drinkers

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Cyprus	544	61.2	48	60.0	592	61.1
Turkey	326	36.7	27	33.8	353	36.4
England	6	0.7	2	2.5	8	0.8
Other	13	1.5	3	3.8	16	1.7
Total	889	100.0	80	100.0	969	100.0

$X^2=5.476$, $df=3$, $p=0.140$, $DA=25(\%2.5)$

There is no statistically significant difference between comparison of birth place distribution of drinkers and non-drinkers according to chi-square statistical method. Mostly the Cypriot students drink more compare with Turkey born students ($X^2=5.476$, $df=3$, $p=0.140$).

Table 41. Comparison of birth place distribution to mothers of drinkers and non-drinkers

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Cyprus	491	55.3	37	46.3	528	54.5
Turkey	375	42.2	39	48.8	414	42.8
England	3	0.3	1	1.3	4	0.4
Other	19	2.1	3	3.8	22	2.3
Total	888	100.0	80	100.0	968	100.0

$X^2=4.145$, $df=3$, $p=0.246$, $DA=26(\%2.6)$

There is no statistically significant difference between comparison of birth place distribution of mothers of drinkers and non-drinkers according to chi-square statistical method. Mothers whose birth place is Turkey and Cyprus drink more than other participants ($X^2=4.145$, $df=3$, $p=0.246$).

Table 42. Comparison of birth place distribution to fathers of drinkers and non-drinkers

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Cyprus	479	53.9	35	43.8	514	53.1
Turkey	384	43.2	42	52.5	426	44.0
England	4	0.5	0	0.0	4	0.4
Other	21	2.4	3	3.8	24	2.5
Total	888	100.0	80	100.0	968	100.0

$X^2=3.793$, $df=3$, $p=0.285$, $CV=26(\%2.6)$

There is statistically significant difference between comparison of birth place distribution of fathers of drinkers and non-drinkers according to chi-square statistical method. Fathers whose birth place is Turkey and Cyprus drink more than other participants ($X^2=3.793$, $df=3$, $p=0.285$).

Table 43. Comparison of distribution to places where drinkers and non-drinkers live

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Village	391	44.0	27	34.2	418	43.2
City	471	53.0	50	63.3	521	53.9
Suburbs	26	2.9	2	2.5	28	2.9
Total	888	100.0	79	100.0	967	100.0

$X^2=3.084$, $df=2$, $p=0.214$, $DA=27(\%2.7)$

There is no statistically significant difference between comparison of distribution of places where drinkers and non-drinkers live according to chi-square statistical method. Participants living in cities drink more than participants living in village and suburbs ($X^2=3.084$, $df=2$, $p=0.214$).

Table 44. Comparison of work status distribution of drinkers and non-drinkers

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Yes	507	57.7	58	73.4	565	59.0
No	371	42.3	21	26.6	392	41.0
Total	878	100.0	79	100.0	957	100.0

$X^2=7.362$, $df=1$, $p=0.007$, $DA=37(\%3.7)$

There is statistically significant difference between comparison of work status distribution of drinkers and non-drinkers according to chi-square statistical method. Compared to unemployed participants, employed participants drink more ($X^2=7.362$, $df=1$, $p=0.007$).

Table 45. Comparison of income status distribution to drinkers and non-drinkers

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Very good	21	2.4	1	1.3	22	2.3
Good	288	32.5	28	35.0	316	32.7
Average	518	58.4	37	46.3	555	57.4
Bad	49	5.5	8	10.0	57	5.9
Very bad	11	1.2	6	7.5	17	1.8
Total	887	100.0	80	100.0	967	100.0

$X^2=75.173$, $df=8$, $p=0.000$, $DA=35(\%35)$

There is statistically significant difference between comparison of income status distribution of drinkers and non-drinkers according to chi-square statistical method. It is seen that participants with average income tend to drink less ($X^2=75.173$, $df=8$, $p=0.000$).

Table 46. Comparison of drinkers and non-drinkers in terms of where they receive most information about drinking

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Media	672	77.7	36	45.6	708	75.0
Book,newspaper,booklet	30	3.5	8	10.1	38	4.0
Family and friends	21	2.4	0	0.0	21	2.2
Friends	49	5.7	20	25.3	69	7.3
Teachers	4	0.5	1	1.3	5	0.5
Doctor, nurse or health officers	10	1.2	0	0.0	10	1.1
Youth centers	0	0.0	1	1.3	1	0.1
Internet	48	5.5	7	8.9	55	5.8
Other	31	3.6	6	7.6	37	3.9
Total	865	100.0	79	100.0	944	100.0

$X^2=75.173$, $df=8$, $p=0.000$, $DA=50(\%5.0)$

There is statistically significant difference between comparison of drinkers and non-drinkers in terms of where they receive most information about drinking according to chi-square statistical method. Although drinkers receive information about drinking mostly from friends but non-drinkers receive most information through media ($X^2=75.173$, $df=8$, $p=0.000$).

Table 47. Comparison of drinkers and non-drinkers in terms of the importance of religion in their lives

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Very important	414	49.0	30	39.5	444	48.2
Relatively important	341	40.4	34	44.7	375	40.7
Unimportant	90	10.7	12	15.8	102	11.1
Total	845	100.0	76	100.0	921	100.0

$X^2=3.302$, $df=2$, $p=0.192$, $DA=73(\%7.3)$

There is statistically significant difference between comparison of drinkers and non-drinkers in terms of the importance of religion in their lives according to chi-square statistical method. Non-drinkers do not pay utmost attention to religion ($X^2=3.302$, $df=2$, $p=0.192$).

Table 48. Comparison of smokers and non-smokers in terms of OPS use

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Non-users of DPM	323	82.2	505	92.3	828	88.1
DPM users	70	17.8	42	7.7	112	11.9
Total	393	100.0	547	100.0	940	100.0

$X^2=22.376$, $df=1$, $p=0.000$, $DA=54(\%5.4)$

There is statistically significant difference between comparison of smokers and non-smokers in terms of OPS use according to chi-square statistical method. OPS use is seen to be higher in smokers ($X^2=22.376$, $df=1$, $p=0.000$).

Table 49. Comparison of smokers and non-smokers in terms of illegal OPS use

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Yes	360	87.6	551	96.8	911	93.0
Non	51	12.4	18	3.2	69	7.0
Total	411	100.0	569	100.0	980	100.0

$X^2=31.16$, $df=1$, $p=0.000$, $DA=14(\%1.4)$

There is statistically significant difference between comparison of smokers and non-smokers in terms of illegal OPS use according to chi-square statistical method. Smokers use illegal OPS four times more than non-smokers ($X^2=31.16$, $df=1$, $p=0.000$).

Table 50. Comparison of smokers and non-smokers in terms of drinking age

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
25 age and below	295	93.1	254	91.0	549	92.1
25 age and above	22	6.9	25	9.0	47	7.9
Total	317	100.0	279	100.0	596	100.0

$X^2=0.834$, $df=1$, $p=0.361$, $DA=398(\%40.0)$

There is no statistically significant difference between comparison of smokers and non-smokers in terms of drinking age according to chi-square statistical method. Smokers aged 25 and below tend to start drinking more ($X^2=0.834$, $df=1$, $p=0.361$).

Table 51. Comparison of smokers and non-smokers in terms of constant drinking age compared to their peers

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Not often	107	87.7	54	85.7	161	87.0
Often	15	12.3	9	14.3	24	13.0
Total	122	100.0	63	100.0	185	100.0

$X^2=0.146$, $df=1$, $p=0.703$, $DA=809(\%81.4)$

There is no statistically significant difference between comparison of smokers and non-smokers in terms of constant drinking age compared to their peers according to chi-square statistical method ($X^2=0.146$, $df=1$, $p=0.703$).

Table 52. Comparison of smokers and non-smokers in terms of the amount of alcohol consumed in one go

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
5 glasses and less	77	18.9	241	43.8	318	33.2
5 glasses of alcohol and more	330	81.1	309	56.2	639	66.8
Total	407	100.0	550	100.0	957	100.0

$$X^2=65,360, df=1, p=0,000, DA=37(\%3,7)$$

There is statistically significant difference between comparison of smokers and non-smokers in terms of the amount of alcohol consumed in one go according to chi-square statistical method. Smokers compared with non-smokers drink 5 glasses of alcohol and more in one go ($X^2=65,360, df=1, p=0,000$).

Table 53. Comparison of smokers and non-smokers in terms of prevalence of drunkenness

Sociodemographic Variables	Smokers		Non-smokers		Total	
	n	%	n	%	n	%
Never drunk	150	37.0	364	65.7	514	53.6
At least once drunk	255	63.0	190	34.3	445	46.4
Total	405	100.0	554	100.0	959	100.0

$$X^2=77.308, df=1, p=0.000, DA=35(\%3.5)$$

There is statistically significant difference between comparison smokers and non-smokers in terms of prevalence of drunkenness according to chi-square statistical method. It's seen that there is a statistical difference. According to this table, it is possible to say that smokers who get drunk at least once smoke more than others ($X^2=77.308, df=1, p=0.000$).

Table 54. Comparison of drinkers and non-drinkers in terms of OPS use

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
No	294	94.5	529	84.8	823	88.0
Yes	17	5.5	95	15.2	112	12.0
Total	311	100.0	624	100.0	935	100.0

$$X^2=18.744, df=1, p=0.000, DA=59(\%5.9)$$

There is statistically significant difference between comparison of drinkers and non-drinkers in terms of OPS use according to chi-square statistical method. Drinkers use OPS three times more than non-drinkers ($X^2=18.744$, $df=1$, $p=0.000$).

Table 55. Comparison of drinkers and non-drinkers in terms of illegal OPS use

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
No	319	97.9	582	90.4	901	92.9
Yes	7	2.1	62	7.1	69	7.1
Total	326	100.0	644	100.0	970	100.0

$X^2=18.328$, $df=1$, $p=0.000$, $DA=24(\%2.4)$

There is statistically significant difference between comparison of drinkers and non-drinkers in terms of illegal OPS use according to chi-square statistical method. It's seen that drinkers use illegal OPS more than non-drinkers ($X^2=18.328$, $df=1$, $p=0.000$).

Table 56. Comparison of drinkers and non-drinkers in terms of smoking cigarette, pipe or cigar

Sociodemographic Variables	Non-drinkers		Drinkers		Total	
	n	%	n	%	n	%
Yes	77	24.2	330	51.6	407	42.5
No	241	75.8	309	48.4	550	57.5
Total	318	100.0	639	100.0	957	100.0

$X^2=65.360$, $df=1$, $p=0.000$, $DA=37(\%3.7)$

There is statistically significant difference between comparison of drinkers and non-drinkers in terms of smoking cigarette, pipe or cigar according to chi-square statistical method. Drinkers smoke cigarette, pipe and cigar significantly more than non-drinkers ($X^2=65.360$, $df=1$, $p=0.000$).

Table 57. Odds ratio and confidence intervals of demographic variables for formation of tobacco use obtained from multivariate logistic regression

Demographic Variables	Tobacco User / Non-user	
	Odds Ratio	%95 CI
Male / Female	1.996	(1.542-2.584)**
25 years old and below / above	1.049	(0.796-1.381)
Living status (lonely / someone)	1.404	(0.918-2.146)
Education level (high school below / and above)	1.149	(0.882-1.497)
Employment (employed / un-employed)	1.810	(1.388-2.360)**
Social Support (no / yes)	1.304	(1.003-1.696)*
Income Level (medium-bad / good)	1.749	(1.083-2.826)*
Living Place (city / village)	1.516	(1.169-1.965)**
Place of birth (Turkey / Cyprus)	1.196	(0.917-1.559)
Alcohol user / non-user	3.552	(2.561-4.925)**
OPS user / non-user	2.606	(1.734-3.916)**
Illicit Drug user / non-user	4.337	(2.493-7.543)**

P<0.05*, P<0.01**, CI=Confidence Interval

Being male, employed, not having social supports, medium-bad income level, living in cities, alcohol use, OPS and Illicit drug use were the risk factors for formation of tobacco use obtained from multivariate logistic regression.

Table 58. Odds ratio and confidence intervals of demographic variables for formation of alcohol use obtained from multivariate logistic regression

Demographic Variables	Alcohol User / Non-user	
	Odds Ratio	%95 CI
Male / Female	3.562	(2.634-4.818)**
25 years old and below / above	1.028	(0.759-1.393)
Living status (lonely / someone)	1.057	(0.658-1.697)
Education level (high school below / and above)	0.411	(0.308-0.549)
Employment (employed / un-employed)	3.153	(2.350-4,228)**
Social Support (no / yes)	0.824	(0.615-1.105)
Income Level (medium-bad / good)	1.038	(0.609-1.770)
Living Place (city / village)	1.501	(1.132-1.990)*
Place of birth (Turkey / Cyprus)	0.428	(0.320-0.571)**
Tobacco user / non-user	3.552	(2.561-4.925)**
OPS user / non-user	3.865	(2.038-7.330)**
Illicit Drug user / non-user	5.332	(2.122-13.395)**

P<0.05*, P<0.01**, CI=Confidence Interval

Being male, employed, living in cities, born in Turkey, tobacco use, OPS and Illicit drug use were the risk factors for formation of alcohol use obtained from multivariate logistic regression.

4. DISCUSSION

This study shows that the rates of cigarette and alcohol use in Cyprus is very high and they are the risk factors for illegal substance use. This study is a continuation study and also comparable with 2003, 2008 and 2013 studies which were used some questionnaire and applied to the same age group. So, it gives information to us about the changes of rates of cigarette and alcohol use in TRNC.

4.1. Cigarette Use

In this study, it is found that at least once in their lifetime cigarette use is %62.7. The lifetime cigarette use was %44.7 in 2003 (Çakıcı et al, 2003), %64 in 2008 (Çakıcı etc. 2014), %62.1 in 2013 (Tutar, 2014). Studies which were conducted in TRNC with high school students show that at least once in lifetime cigarette use is between %26.8 and %47.2 (Çakıcı ve Çakıcı, 1996; 1999; Çakıcı et al, 2000; Eş, 2011). In a study which was conducted in 2012 with university students it was found that at least once in their life time cigarette use was %69.5 (Çakıcı et al, 2010). Those results show that cigarette use increase from adolescence to adulthood. The current prevalence of cigarette use in TRNC is %41.8. According to the data of World Bank, the prevalence of smoking cigarettes in East Asia and Pacific is %34, in Europe and Middle Asia is %35, in Latin America and Caribbean is %32, in Middle East and North Africa is %21, in South Asia is %20 (Anderson, 2006). It can be said that the higher prevalences of cigarette smoking were seen in Europe and Middle Asia countries. The prevalence of cigarette use in TRNC is higher than average rates of its own geographic location Europe and also higher than the average rates from all over the areas of world. Smoking cigarette rate is higher in TRNC according to the studies that were conducted in 50 states of America, Colombia in 2009-2010 (King et al, 2012) and in Tahrán (Fotouhi et al, 2009). The rates in TRNC are similar like El Salvador (%42.7), Guatemala (%43.1) and Honduras (%43.8) Latin America countries (Tong et al, 2011). However, the rates in Ukraine (%66.8), Russia (%63) and Turkey (%60.3) are higher than the results of our study (Ögel et al 2003; Bobak et al, 2006). The main reasons of the high cigarette use rates are seen in TRNC. Acceptability of cigarette use are high in TRNC society, the lack of adequate legislation and supervision, the widespread availability of cigarette advertising and low price of cigarette. TRNC also is a touristic island that can cause high alcohol use in

entertainment places accompany with smoking cigarette. In this study, it was found that men that live alone and 18-25 ages smoke more cigarettes. Moreover, smoking cigarette rates are higher in cities. The conditions of city life make people to smoke more cigarettes (Fetihi, 2002). Working people smoke more cigarettes. Work stress and the conditions of work are effective on smoking cigarettes (Şahin et al., 2007). Economic conditions are also effective on smoking cigarettes. There is not significant difference between people who smoke cigarettes and do not smoke according to religion. In other studies which were conducted in TRNC showed that religion does not have any effect on smoking cigarettes (Akfert et al., 2009).

Being male, employed, not having social supports, medium-bad income level, living in cities were the risk factors for formation of tobacco use. It's found that there is a close relationship between alcohol use and tobacco use. In the study of which is conducted by Akfert et al., 2009, It's found there is strong relationship between tobacco use and alcohol use. In 2004 the study which is conducted by Karatay and Kubilay (2004), it's found there is a high correlation between tobacco use and alcohol use. People who smoke cigarettes are more prone to be drunk 5 and more glasses of alcohol in one time. Both on the other hand and illegal substance use are higher among people who smoke cigarettes. Cigarette use is a risk factor for OPS use and there is relationship between beginning cigarette in early ages and OPS use (Çakıcı et al., 2014). Studies showed that there is close relation between OPS use and tobacco use. Sex, education level of father, family communication, cigarette and alcohol are the risk factors for substance use (Ulukoca et al., 2013). While examining having dependent friends whether or not change the tobacco use behavior of students, it has been found that students who smoke have %37.6 dependent friends. Students who do not smoke have % 17.9 dependent friends (Erdamar and Kurupinar, 2014). It has been seen that there is significant relationship between having dependent friend and tobacco use. (Erdamar and Kurupinar, 2014).

4.2. Alcohol Use

In our study at least once in their life time alcohol use is found as 72.1%. At least once in their lifetime alcohol use is found as % 82.1 in 2003, %77.1 in 2008, and %68.5 in 2013. When the datas of 2013 and 2015 are compared, it is found that

alcohol use increased. At least once in their lifetime alcohol use in high school study is found as %85.9 (Çakıcı et al, 2010) and in university it is found as %81.0 (Çakıcı et al, 2014). In a study which is conducted in Istanbul in 15 different studies, it is found that prevalence of alcohol use is %51.2 (Ögel et al, 2006), and it is found in university students as in between 30-76% (Altındağ et al, 2005; Yılmaz et al, 2007). In a study in which the rates of alcohol use in Australia and USA 1995, 1998, 2001, 2002 and 2004 are compared, it is found that at least once in their life time alcohol use in Australia 87.8-90.4, in USA is 83.7-84.8% (Maxwell et al, 2006). The prevalence of alcohol use in TRNC is similar with Australia and USA. However, when the prevalence of alcohol use of TRNC and Turkey is compared, it is found that (Çakıcı et al, 2014), TRNC has higher prevalence of alcohol use (World Health Organization, 2004). Cyprus is a touristic island, it has free advertisements, legal procedures are not enough, so they lead to higher alcohol use rates. Furthermore, universities have younger populations, alcohol can be accessed easily, advertisements, lower prices and cultural factors lead to increase in alcohol use (Çakıcı et al, 2003; Çakıcı et al, 2010).

In this study, the men who are 18-25 years and live alone or with their parents have high alcohol use. People who have level education use more alcohol than people who have lower education level. Education level do not prevent alcohol use. Çakıcı et al., 2014, declared that for prevention of alcohol use, special education programs are necessary and ordinary education is not enough to prevent alcohol use. Moreover, alcohol use is higher among people who live in cities. It can be said that there is positive relationship between higher alcohol use rates and to be drunk and smoking cigarettes. People who use alcohol have 3 more times other psychoactive substance than people who do not use alcohol. Moreover, people who use alcohol smoke more cigarettes, pipe and cigar.

The study It observed that there is relationship between other psychoactive substance and alcohol use. While examining having dependent friends whether or not change the alcohol use behavior of students, it has been found that students who smoke have %37.7 dependent friends, students who do not smoke have % 19.1 dependent friends. Erdamar and Kurupınar (2014) declared that there is significant relationship between having dependent friend and alcohol use.

5. CONCLUSION

This study shows that cigarettes and alcohol are the most acceptable and used psychoactive substances. In TRNC society, cigarette and alcohol use effect each other. Cigarettes and alcohol are more common in society and they become the risk factors for TRNC culture. This study also shows that cigarettes and alcohol use are related with other psycgoactive substance and illegal substance use. They are also the risk factors for OPS use. Cigarettes and alcohol prevention programs are necessary to prevent OPS use. So, prevention programs to prevent OPS are not enough. Education programs in primary school to prevent alcohol and cigarettes use can be effective to decrease alcohol and cigarettes use and also OPS use. As a result, in TRNC multidiciplinary approach is needed and also public health policies are necessary for prevention of cigarettes and alcohol use.

REFERENCES

Adelson, M. Betty, Paul Casadonte, Mary Jeanne Kreek, Orna Levran, Stephanie Linzy, David Londono, Ann Nielsen, Kieron O'Hara, Jonathan Ott, Elior Peles, Mike Randesi, Rotrosen John, 2008. Genetic susceptibility to heroin addiction: A candidate gene association study. **Genes, Brain, and Behavior**, v. 7. n: 720-729.

Agrawal, Arpana, Hinrichs . Anthony L, Gerald Dunn, Sarah Bertelsen, Danielle M. Dick, Scott F. Saccone, Nancy L Saccone, Richard A. Grucza, Jen C. Wang, C. Robert Cloninger, Howard J. Edenberg, Tatiana Foroud, Victor Hesselbrock, John Kramer, Kathleen K. Bucholz, Samuel Kuperman, John I. Numberger Jr, Bernice Porjesz, Marc A. Schuckit, Alison M. Goate, Laura J. Bierut, 2008. Linkage scan for quantitative traits identifies new regions of interest for substance dependence in the Collaborative Study on the Genetics of Alcoholism (COGA) sample. **Drug and Alcohol Dependence**. v. 93. n.1: 12-20.

Agrawal, Arpana, Melanie Lynskey, 2008. Are there genetic influences on addiction:evidence from family, adoption, and twin studies. **Addiction**. v. 103. n.7: 1069-1081.

Allgulander, Christer, 1989. Psychoactive drug use in a general population sample, Sweden: correlates with perceived health, psychiatric diagnoses, and mortality in an automated record-linkage study. **American Journal of Public Health**. v. 79. n. 8: 1006-1010.

Barış, İzzettin. 1994. **Sigara ve Sağlık**. İstanbul: MEB Yayinlari. 2464.

Barut, Yasar, 1992. Üniversite Öğrencisinin Sigara, Alkol, İlaç Alma ve Uyuşturucu Madde Kullanım Alışkanlığı, **Araştırma Sağlık Dergisi**. v. 64. n. 1: 11-16.

Bayar, Reha, Mesut Yavuz. 2008. **Alkol Bağımlılığı**, İ.Ü. Cerrahpaşa Tıp Fakültesi Sürekli Tıp Eğitimi Etkinlikleri, Sempozyum Dizisi No:62 Mart 2008.

Bernat, Edward, Hicks Bill, Iacono William, Krueger Robert, Malone Steve, McGue Matthew, Patrick Coggins. 2007. **Genes med iate the association between P3 amplitude and externalizingdisorders**. Psychophysiology.

Bilir, Nazmi, Hilal Özcebe, Dilek Aslan, Toker Ergüder. Yardım Mahmut Sadi, Eser Sultan, Telatar Gökhan, 2007. **Avrupa Tütün Kontrolü Raporu 2007-Türkçe** Çevirisi. Bilir, N. Ed. WHO Yayınları, 2008, 17.

Brown, Stephanie, Irvin Yalom, 2008. **Alkolizm terapisi**. (Çev. T. Kazmirci). İstanbul: Prestij Yayınları.

Chen, An, Anh Ho, Yuferov Ji, Mike Kreek, Orna Levran, David Nielsen, Jonathan Ott, 2008. Genotype patterns that contribute to increased risk for or protection from developing heroin addiction. **Molecular Psychiatry**. v.7: 417-428.

Connors, Gerard, Stephen Maisto, David Watson. 1988. Racial factors influencing collegestudents' ratings of alcohol's usefulness. **Drug Alcohol Dependend**. v.21: 247-52.

Çakıcı, Mehmet, 1999. Madde Kullanıcılarının Özellikleri: Türkiye'de Çok Merkezli Bir Araştırma (Birinci Aşama). **3P Dergisi**. v. 7. n. 1: 34-36.

Çakıcı, Mehmet. 2000. **KKTC Araştırmaları ile Uyuşturucu Madde Kullanımı**. Lefkoşa: KKTC Devlet Basımevi.

Çakıcı, Mehmet, Ebru Çakıcı. 2000a. **K.K.T.C. Lise Gençliğinde Uyuşturucu**

Madde Kullanım Yaygınlığı, 1996. Lefkoşa: KKTC Başbakanlık Devlet Basımevi.

Çakıcı, Mehmet, Ebru Çakıcı. 2000b. **K.K.T.C. Lise Gençliğinde Uyuşturucu**

Madde Kullanım Yaygınlığı, 1999. Lefkoşa: KKTC Başbakanlık Devlet Basımevi.

Çakıcı, Mehmet, Ebru Çakıcı, Bingül Subaşı. 2001. **KKTC Ortaokul Öğrencileri arasında Madde Kullanım Yaygınlığı. 8. Ulusal Sosyal Psikiyatri Kongresi Bilimsel Çalışmaları** (28-31 Mayıs 2001, Lefkoşa). 176-182. Lefkoşa.

Çakıcı, Mehmet, Ebru Çakıcı, Zafer Bekiroğulları, et al. 2003. **Kuzey Kıbrıs'ta**

Madde Kullanımının Yaygınlığı. Lefkoşa: KKTC Devlet Basımevi.

Çakıcı, Mehmet, Ebru Çakıcı, Ayhan Eş. 2010. Kuzey Kıbrıs Türk Cumhuriyeti Lise Öğrencilerinde Psikoaktif Maddeleri Kullanma Yaygınlığı. **Anatolian Journal of Psychiatry**. v. 11: 206-212.

Çakıcı, Mehmet, Ebru Çakıcı, Meryem Karaaziz, Neslihan Tutar, Ayhan

Eş, 2014. KKTC'de Psikoaktif Maddelerin Kullanım Yaygınlıkları ve Risk

Etkenleri. **Bağımlılık Dergisi**. v. 15. n.4: 159-166.

Çakıcı Ebru, Mehmet Çakıcı, Ayhan Eş, Deniz Ergün. 2014. The prevalence and risk factors of substance use among university students in Turkish Republic of Northern Cyprus. **Anatolian Journal of Psychiatry**. v. 15. n.2: 108-115.

Çivi Serhat, Kemal Şahin. 1991. Selçuk Üniversitesi Tıp Fakültesi ve Sağlık Hizmetleri Meslek Yüksekokulu öğrencilerinin sigara konusundaki bilgi tutum ve davranışları. **Aile ve Toplum**

Dağlı, Elif. 1994. **Sigara Bağımlılığı ve Nedenleri; Sigara ve Sağlık**. MEB Yay: 2464. İstanbul. S.16-17.

Derdiman, Ramazan Cengiz. 2006. Uyuşturucu Psikoaktif psikoaktif maddeler Hakkında Genel Bilgi ve Uyuşturucu Psikoaktif psikoaktif maddelerle Mücadele Esasları. **Akademik Araştırmalar Dergisi**. v.28: 103-122.

Devlet Denetleme Kurumu, Araştırma ve İnceleme Raporu, 2014. **Madde ve Diğer Bağımlılıklar ile Mücadele Kapasitesinin ve Bu Bağlamda Türkiye Yeşilay Cemiyetinin Değerlendirilmesi**.

Erdamar, Gürcü, Abdülhamit Kurupınar. 2014. Ortaöğretim Öğrencilerinde Görülen Madde Bağımlılığı Alışkanlığı ve Yaygınlığı: Bartın İli Örneği, **Sosyal Bilimler Dergisi**. v.16. n. 1.

Esirgemez, Cihan, Çakıcı Mehmet. 2014. Evaluation Of The Cannabis As A Gateway Drug: Gateway Hypothesis. Yüksek lisans tezi. Yakın Doğu Üniversitesi. Sosyal Bilimler Enstitüsü. Lefkoşa.

Haley, George, David Hoffmann. 1985. Analysis for nicotine and cotinine in hair to determine cigarette smoker status. **Clin Chem**. v. 31. 1598-1600.

Hasin, S. Deborah, Frederick Stinsons, Elizabeth Ogburn, Grant F, Bridget. 2007. Prevalence, correlates, disability, and comorbidity of DSM-IV alcohol abuse and dependence in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. **Archives of general psychiatry**. v. 64(7). n: 830-842.

Hawkins, J. David, Catalano Richard, Miller Janet, 1992. Risk And Preventive Factors For Alcohol And Other Drug Problems In Adolescence And Early Adulthood: Implications For Substance Abuse Prevention. **Psychol Bull.** v. 112. n. 1: 64-105

Hogan, Michael J. 2000. Diagnosis And Treatment Of Teen Drug Use. **Med. Clin. North Am.** v. 84: 927-66.

Jellinek, Elvin, Morton. 1952. Phases of alcohol addiction. **Quarterly journal of studies on alcohol.** v. 13. n:4. 673-684.

Kalyoncu Aytaş, Hasan Mırsal. 2000. Alkol kullanım bozuklukları. **Psikiyatri Dünyası.** 2000.

Karatay, Gülnaz, Gülümser Kubilay. 2004. Sosyo ekonomik Düzeyi Farklı İki Lisede Madde Kullanımı ve Etkileyen Faktörlerin Belirlenmesi. Hacettepe Üniversitesi Sağlık Bilimleri Enstitüsü. Bilim Uzmanlığı tezi. **Hemşirelikte Araştırma Geliştirme Dergisi.** v. 6. n. 1-2: 57-70.

Kaya, Sefa. 1991. Sigara ve Sağlığımız, **Bilim ve Teknik.** v. 24. n. 289: 42-45, Aralık 1991.

Keskinoglu, Pembe, Neslihan Karakuş, Metin Pıçakçıefe ve ark. 2006. İzmir'de lise öğrencilerinde sigara içme sıklığı ve içicilik davranışı üzerine sosyal öğrenmenin etkisi. **Toraks Dergisi.** v.7. n: 190-5.

Köknel, Özcan. 1998. **Bağımlılık.** İstanbul: Altın Kitaplar Yayınevi.

Kuppin, Sara, Richard Carpiano. 2006. Public conceptions of serious mental illness and substance abuse, their causes, and treatments: Findings from the 1996 General Social Survey. **American Journal of Public Health**. v. 96. n: 1766-1771.

Mangır, Mustafa, Neriman Aral, Gülen Baran. 1992. Üniversite Öğrencilerinin Alkol Kullanımlarının İncelenmesi. **Gıda Dergisi**, v.17. n:5.

McKee Sally, David R Hinson, Rounsaville Paul D, John Petrelli, 2004. Survey of subjective effects of smoking while drinking among college students. **Nicotine & Tobacco Research**. 2004. v.6. n.1: 111-117.

Ögel, Kültegin. 1997. **Uyuşturucu Psikoaktif psikoaktif maddeler ve Bağımlılık**. İstanbul: İletişim Yayınları.

Runeson, Bo. 1990. Psychoactive substance use disorder in youth suicide. **Alcohol and Alcoholism**. v. 25. n. 5: 561-568.

Sağlık Bakanlığı. 1995. Lise öğrencilerinin sigara, alkol ve uyuşturucu maddeler konusundaki bilgi tutum ve davranışlarının belirlenmesi projesi. **Temel Sağlık Hizmetleri Genel Müdürlüğü Ruh Sağlığı Daire Başkanlığı**.

Sezer, Refik. 1984. **Ankara'nın Ortaokul, Lise ve Lise Dengi Okullarında Sigara İçeme Durumu ve Sigaraya Başlama Nedeni**. Tübitak tarafından desteklenen Araştırma raporu.

Schuckit, Marc. 2000. **Alcohol-Related disorders**. Sadock B.J. Sadock V.A. Ed. Kaplan. Sadock's comprehensive textbook of psychiatry. 2008. Philadelphia: Lippincott Williams. Wilkins.

Tanrikulu, A. Çetin, Çarman Kürşat Bora, Palancı Yılmaz, Çetin Didem, Karaca Mehmet. 2009. Kars il merkezinde çeşitli üniversite öğrencileri arasında sigara kullanım sıklığı ve risk faktörleri. **Türk Toraks Dergisi**. v. 10, n: 101-106.

The Model European Questionnaire Survey Handbook. 2001-2002. **The initial inventory of surveys including questions on drug availability**: Lizbon: EMCDDA.

Tol, Canan, 1990. **Lise öğrencilerinin alkol kullanımını etkileyen bazı değişkenler**. Unpublished Master Thesis. Hacettepe University. Ankara.

Turhan, Esra, Tacettin İnandı, Cahit Özer, Sabahat Akoğlu. 2011. Üniversite Öğrencilerinde madde kullanımı, siddet ve bazı psikolojik özellikler. **Türkiye Halk Sağlığı Dergisi**, v. 9(1). n: 33.

Tütar, Neslihan. 2013. The Prevalence of Substance Use in Northern Cyprus. **Unpublished Master Thesis**. Near East University Graduate School of Social Sciences.

Ulukoca, Neşe, Şenlen Gokgoz, Ali Karakoç. 2013. Kırklareli Üniversitesi Öğrencileri Arasında Sigara, Alkol ve Madde Kullanım Sıklığı, Kırklareli Üniversitesi Sağlık Hizmetleri MYO. *Firat Med J*. V. 18. n. 4: 230-234

WHO. 1993. **The Health of Young People: A Challenge and a Promise**, Geneva: World Health Organization.

Yoshimoto, K, McBride, W. J, Lumeng, L, Li, T. K. 1992. Alcohol stimulates the release of dopamine and serotonin in the nucleus accumbens. **Alcohol**. v. 9(1). n: 17-22.

APPENDIXES

Appendix 1. Informed Consent Form

AYDINLATILMIŞ ONAM FORMU

Bu çalışma, Yakın Doğu Üniversitesi Fen Edebiyat Fakültesi Psikoloji Bölümü tarafından gerçekleştirilen bir çalışmadır.

Bu çalışmanın amacı KKTC'deki madde kullanım yaygınlığını araştırmak, sorunun boyutlarını öğrenmek ve giderek yükselen madde bağımlılığı sorununun nedenlerini araştırmaktır. Çalışma sonucunda elde edilen veriler doğrultusunda toplum genelinde madde kullanımını önlemeye yönelik bilimsel programların geliştirilmesi amaçlanmaktadır.

Anket tamamen bilimsel amaçlarla düzenlenmiştir. Anket formunda kimlik bilgileriniz yer almayacaktır. Size ait bilgiler kesinlikle gizli tutulacaktır. Çalışmadan elde edilen veriler yalnızca istatistik veri olarak kullanılacaktır. Yanıtlarınızı içten ve doğru olarak vermeniz bu anket sonuçlarının toplum için yararlı bir bilgi olarak kullanılmasını sağlayacaktır.

Telefon numaranız anketörün denetlemesi ve anketin uygulandığının belirlenmesi amacıyla istenmektedir.

Yardımanız için çok teşekkür ederim.

Uzman Psikolog,
Meryem Karaaziz.

Yukardaki bilgileri ayrıntılı biçimde tümünü okudum ve anketin uygulanmasını onayladım.

İsim:

İmza:

Telefon:

BİLGİLENDİRME FORMU

KUZEY KIBRIS TÜRK CUMHURİYETİNDE MADDE KULLANIMININ YAYGINLIĞI, 2015

Bu çalışmanın amacı KKTC'deki madde kullanım yaygınlığını araştırmak, sorunun boyutlarını öğrenmek ve giderek yükselen madde bağımlılığı sorununun nedenlerini araştırmaktır. Çalışma sonucunda elde edilen veriler doğrultusunda toplum genelinde madde kullanımını önlemeye yönelik bilimsel programların geliştirilmesi amaçlanmaktadır..

Bu çalışmada size bir demografik bilgi formu ve bir dizi ölçek sunduk. Demografik bilgi formu sizin yaş cinsiyet gibi demografik özellikleriniz hakkındaki soruları içermektedir. Ölçekler ise madde kullanım sorunun düzeyini, risklerini ve madde kullanım davranışının özelliklerini ölçmektedir.

Daha önce de belirtildiği gibi, ölçeklerde ve görüşmelerde verdiğiniz cevaplar kesinlikle gizli kalacaktır. Eğer çalışmayla ilgili herhangi bir şikayet, görüş veya sorunuz varsa bu çalışmanın araştırmacılarından biri olan Uzm. Psk. Meryem Karaaziz'le iletişime geçmekten lütfen çekinmeyiniz (meryem.karaaziz@yahoo.com, telefon: 0392 22 36 464) (iç hat: 254).

Eğer bu çalışmaya katılmak sizde belirli düzeyde stres yaratmışsa ve bir danışmanla konuşmak istiyorsanız, askeriyede ücretsiz hizmet veren askeri psikoloğa başvurabilirsiniz.

Eğer araştırmanın sonuçlarıyla ilgileniyorsanız, Haziran 2015 tarihinden itibaren araştırmacıyla iletişime geçebilirsiniz.

Katıldığınız için tekrar teşekkür ederim.

Uzman Psikolog,

Meryem Karaaziz

Psikoloji Bölümü,

Yakın Doğu Üniversitesi,

Lefkoşa.

Appendix 2. Demographic Information Form

SOSYAL SORUNLAR ANKETİ

Bu anket çalışması sosyal sorunlarımızı ve alışkanlıklarımızı araştırmaya yönelik bilimsel bir çalışmamızdır. Kıbrıs genelinde 18-65 yaş grubundaki kadın-erkek bireylere uygulanacaktır. Bu çalışmada kesinlikle kimlik bilgileri kullanılmayacaktır.

Yalnızca çalışmanın istatistik verileri bilimsel olarak akademisyenler tarafından ülkemizdeki sorunların çözümüne yönelik kullanılacaktır.

Katkı sağladığınız için teşekkür ederiz.

BÖLÜM I

Kendiniz ve Aileniz Hakkında

1.Cinsiyetiniz nedir?

1.Erkek 2 .Kız

2.Son doğum gününüzde kaç yaşınızı doldurdunuz?

3.Halen kimlerle yaşıyorsunuz?

1.Yalnız 2.Eşimle-Partnerimle 3.Çocuklarımla 4.Anne ve babamla
5.Anne veya babamla
6.Üvey anne veya üvey babamla 7.Akrabalarımla 8.Arkadaşlarla 9.Diğer

4a.Ulaşılan öğretim düzeyi

1.Okula Gitmedim 2.İlkokul 3.Ortaokul 4.Lise 5. Üniversite ve üzeri

4b-c. Anne-babanızın ulaştığı öğrenim düzeyi

	Baba	Anne
1. Okula Gitmedim		
2. İlkokul		
3. Ortaokul		
4. Lise		
5. Üniversite ve üzeri		

4d. Doğum yeriniz neresidir?

1. Kıbrıs 2. Türkiye 3. İngiltere 4. Diğer

4e. Annenizin doğum yeri neresidir?

1. Kıbrıs 2. Türkiye 3. İngiltere 4. Diğer

4f. Babanızın doğum yeri neresidir?

1. Kıbrıs 2. Türkiye 3. İngiltere 4. Diğer

5. Yaşamınızın büyük çoğunluğunu nerede geçirdiniz?

1. Köy 2. Şehir 3. Şehir dış mahallesi

6. Çalışıyor musunuz ? 1. Evet 2. Hayır

7. Sosyal (Devlet, aile, arkadaş maddi ve manevi desteği) desteğiniz var mı? 1.

- Evet 2. Hayır

8. Gelir durumunuz nasıldır? 1. Çok iyi 2. İyi 3. Orta 4. Kötü

5. Çok kötü

BÖLÜM II

Bilgi kaynakları

1. Uyğururucu maddeler hakkında bilgi en fazla nereden duyduğunuzu daire içine alınız.

(sadece birini)

1. Basından
2. Kitap, dergi, broşür vb.
3. Aile ve akrbalar
4. Arkadaşlar
5. Okul Öğretmenleri
6. Doktor, hemşire veya sağlık görevlisi
7. Gençlik merkezi
8. İnternet
9. Diğer

BÖLÜM III

Uyğururucu maddeler hakkında bilgi

1.Din hayatınızda ne kadar önemlidir?

- 1.Çok önemli 2.Kısmen önemli 3.Önemli değil

2-13. Aşağıdaki maddeleri hiç duydunuz mu?

	Hayır	Evet
2. Uçucu Madde (Tiner, bali vs)	1	2
3. Yatıştırıcı (Valium, Diazem)	1	2
4. Esrar	1	2
5. Bonzai	1	2
6. Amfetamin	1	2
7.Ecstasy	1	2

8. Kokain	1	2
9. Relevin	1	2
10. Eroin	1	2
11. LSD	1	2
12. Kodeinli Şurup	1	2
13. Hap (akineton, roş-rohypnol, nembutal-sarı bomb vs.)	1	2
14. Anabolizan Steroid	1	2

BÖLÜM IV

Boş zaman etkinlikleri

14.Boş zamanınızın çoğunu kiminle geçirirsiniz? (sadece bir cevap işaretleyiniz.)

- | | | | |
|--------------------|--------------|-----------------|----------|
| 1. Yalnız | 2. Anne-baba | 3. Arkadaşlar | 4. |
| Eş/Sevgili/partner | | | |
| 5.Kardeşlerimle | 6. Akrabalar | 7. Başkalarıyla | 8. Diğer |

15.Okul dönemi boyunca boş zamanınızın çoğunu nerede geçirirsiniz? (sadece bir cevap işaretleyiniz.)

- | | |
|--------------------|---------------------------------|
| 1.Kendi evimde | 6.Gençlik kulübü/ spor klubünde |
| 2.Akrabamın evinde | 7.Club/Diskotek/ dans salonunda |
| 3.Arkadaşın evinde | 8.İnternet kafede |
| 4.Caddede/ sokakta | 9.Kafe/restorant |
| 5.İş yerinde | 10.Kahvede |
| | 11.Diğer |

BÖLÜM V

Sağlık davranışları hakkında görüşler

Kişilerin aşağıdaki etkinliklere katılmasını onaylayıp onaylamadığınızı lütfen belirtiniz. (Her biri için tek rakam işaretleyiniz.)

	Kesinlikle Onaylarım	Onaylarım	Kararsızım	Onaylamam	Kesinlikle Onaylamam
1.Sigara içmek	1	2	3	4	5
2.Bira, şarap gibi alkollü içecekler içmek	1	2	3	4	5
3.Viski, votka, konyak gibi daha sert içkiler içmek	1	2	3	4	5
4.Bar veya meyhaneye gitmek	1	2	3	4	5
5.Marihuana veya haşış (esrar, ot) kullanmak	1	2	3	4	5
6.Bonzai içmek	1	2	3	4	5
7. Eroin veya kokain benzeri uyuşturucu madde kullanmak	1	2	3	4	5

8. Doktor reçetesi olmadan 1 2 3 4 5
sakinleştirici-uyku ilacı
almak

BÖLÜM VI

Alkol ve Uyuşturucu Madde Kullanımı

Bu bölümde size bazı kişisel sorular sormak ve sizin için hangilerinin geçerli olduğunu öğrenmeyi arzu ediyoruz. Bu soru formunun tamamen anonim olduğunu (kimlik bilginizi içermediğini) hatırlatarak, mümkün olduğunca dürüst cevaplamanızı rica ediyoruz.

1. Eğer bir uyuşturucu madde kullandıysanız bu nerede oldu?

1 .Hiç kullanmadım 2. Kıbrıs'da 3.Yurtdışında

2. Eğer herhangi bir uyuşturucu madde kullandıysanız, kullanma nedeniniz neydi?

(sadece bir cevap işaretleyiniz)

1.Hiç kullanmadım	4.Merak	7.Tepki-Kızgınlık
2. Arkadaşlar içtiği için	5. Yalnızlık	8.Düşüncesizlik
3 .Bir ilişkiyi sürdürebilmek	6 Başkasının baskısı	9.
Başka(.....)		

3. Trafik kurallarına uymama nedeniyle hiç polisle sorun yaşadınız mı?

1.Hayır 2.Evet

4. Yasadışı herhangi bir eylem nedeniyle hiç polisle sorun yaşadınız mı?

1.Hayır 2.Evet

5. Uyuşturucu madde ile ilgili olarak hiç polisle sorun yaşadınız mı?

1.Hayır 2.Evet

6. Arkadaşlarınız hiç yasadışı bir etkinlikte bulundu mu?

1. Hayır 2. Evet

7. Uyuşturucu madde kullanmaya karar verseniz, nereden bulacağınızı biliyor musunuz?

1. Hayır 2. Evet

8. Alkol etkisindeyken kaç defa cinsel ilişki yaşadınız?

1. Hiçbir zaman 2. 1-2 kere 3. 3 veya daha fazla 4. Her zaman

9. Esrar, eroin, kokain gibi bir uyuşturucu madde etkisinde kaç defa cinsel ilişki yaşadınız?

1. Hiçbir zaman 2. 1-2 kere 3. 3 veya daha fazla 4. Her zaman

10. Eğer Kıbrıs'da bir uyuşturucu madde kullandıysanız, bu nerede oldu?

- | | |
|----------------------------|--|
| 1. Evde tek başıma | 6. Gençlerin bulunduğu başka bir yerde |
| 2. Evde arkadaşlarımla | 7. Okulda |
| 3. Arkadaşımın evinde | 8. Askerde |
| 4. Caddede/ sokakta | 9. Başka bir yerde |
| (.....) | |
| 5. Bir meyhane veya klüpte | 10. Hiç uyuşturucu almadım |

11. Evinizde alkol en çok ne zaman servis edilir? (sadece bir cevap işaretleyiniz)

- | | |
|--------------------------------------|-------------------------|
| 1. Hiçbir zaman | 5. Öğle yemeğinde |
| 2. Misafirimiz olduğun | 6. Akşam yemeğinde |
| 3. Herhangi bir zaman, meze olmadan, | 7. Sadece Pazar günleri |
| rahatlatıcı bir içecek olarak | 8. Bir şeyler kutlarken |
| 4. Herhangi bir zaman, mezeyle | 9. Diğer (.....) |

12. Kaç yaşında uyuşturucu ilacı almaya başladınız? (eğer bu olduysa)
(.....)

BÖLÜM VII

Tütün ve Alkol

1- Sigara, pipo ya da puro gibi tütün içiyor musunuz ☐1 Evet ☐2 Hayır

2-Hayatınız boyunca kaç kez sigara içtiniz?

1)☐02)☐1-2 3) ☐3-54)☐6-95) ☐10-19 6)☐20-39 7)☐40-veya daha fazla

3-Son 12 ayda kaç kez sigara içtiniz?

1) ☐0 2) ☐1-2 3) ☐3-5 4) ☐6-9 5) ☐10-19 6) ☐20-39 7) ☐40-veya daha fazla

4- Son 30 günde ne sıklıkla sigara içtiniz?

- | | | | |
|---------------------------|--------------------------|----------------------------|--------------------------|
| 1- Hiç içmedim | <input type="checkbox"/> | 6-Günde 11-20 sigara | <input type="checkbox"/> |
| 2- Haftada 1 sigaradan az | <input type="checkbox"/> | 7-Günde 20-30 sigara | <input type="checkbox"/> |
| 3- Günde 1 sigaradan az | <input type="checkbox"/> | 8-Günde 31-40 sigara | <input type="checkbox"/> |
| 4- Günde 1-5 sigara | <input type="checkbox"/> | 9-Günde 40 sigaradan fazla | <input type="checkbox"/> |
| 5- Günde 6-10 sigara | <input type="checkbox"/> | | |

5- Eğer sigara kullanıyorsanız, hiç sigarayı bırakmakta zorlandınız mı?

1)☐Çok zorlandım 2)☐Zorlandım 3)☐Zorlanmadım 4)☐Hiç zorlanmadım

6- Kaç yaşında sigara almaya başladınız? (eğer bu olduysa)

(.....)

7- Kaç yaşında alkol almaya başladınız? (eğer bu olduysa)

(.....)

8- Kaç yaşında sürekli olarak yaşitlarımızdan daha fazla miktarda alkol kullanmaya başladınız?
(eğer bu olduysa) (.....)

9-Hayatınız boyunca kaç kez alkollü bir içecek içtiniz?

- 1) ☐ 0 2) ☐ 1-2 3) ☐ 3-5 4) ☐ 6-9 5) ☐ 10-19 6) ☐ 20-39 7) ☐ 40-veya daha fazla

10-Son 12 ay içinde kaç kez alkollü bir içecek içtiniz?

- 1) ☐ 0 2) ☐ 1-2 3) ☐ 3-5 4) ☐ 6-9 5) ☐ 10-19 6) ☐ 20-39 7) ☐ 40-veya daha fazla

11-Son 30 gün içinde kaç kez alkollü bir içecek içtiniz?

- 1) ☐ 0 2) ☐ 1-2 3) ☐ 3-5 4) ☐ 6-9 5) ☐ 10-19 6) ☐ 20-39 7) ☐ 40-veya daha fazla

12- Son 30 gün içinde alkol kullanma sıklığınız ne kadardır?

- | | | |
|------------------------|-------------------|-------------------|
| 1- Hiç | 3-İki haftada bir | 5-Günde bir kez |
| 2-Haftada ikiden fazla | 4-Haftada bir | 6-Haftada iki kez |

13- Bir seferde genellikle ne kadar alkol alırsınız?(bir içki, bir şişe veya bardak bira, bir kadeh şarap, bir bardak rakı veya diğer içkiler anlamında kuhanılmaktadır.)

- 1) ☐ Alkol içmem 2) ☐ 1-2 içki 3) ☐ 3-4 içki 4) ☐ 5 veya fazla içki

14-En son içki içtiğinizde neredeydiniz;

- | | |
|------------------------------|---|
| 1-Hiç bir zaman içki içmem | 2-Evdeydim |
| 3-Başka birisinin evindeydim | 4-Dışarda, sokak, park veya açık havadaydım |
| 5-Bir barda veya kafedeydim | 6-Diskodaydım |
| 7-Lokantadaydım | 8-Diğer(lütfen belirtiniz) |

15-Hayatınız boyunca kaç kez içki içtiğiniz için sarhoş oldunuz?

1) ☐0 2) ☐1-2 3) ☐3-5 4) ☐6-9 5) ☐10-19 6) ☐20-39 7) ☐40-veya daha fazla

16-Son 12 ay içinde kaç kez içki içtiğiniz için sarhoş oldunuz?

1) ☐0 2) ☐1-2 3) ☐3-5 4) ☐6-9 5) ☐10-19 6) ☐20-39 7) ☐40-veya daha fazla

17-Son 30 gün içinde kaç kez içki içtiğiniz için sarhoş oldunuz?

1) ☐0 2) ☐1-2 3) ☐3-5 4) ☐6-9 5) ☐10-19 6) ☐20-39 7) ☐40-veya daha fazla

18-Hangi nedenlerden dolayı alkollü içki kullanıyorsunuz?(Birden fazla yanıt işaretleyebilirsiniz)

- | | | |
|----------------|-----------------------|-----------------------------|
| 1. Eğlenmek | 4. Sinirlendiğim için | 7. Sorunlarımdan uzaklaşmak |
| 2. Uyuyabilmek | 5. Sıkıntıdan | 8. Arkadaşlarım |
| 3. Denemek | 6. Rahatlamak | 9. Kendimi iyi hissetmek |

YATİŞTİRİCİ

1. Yatıştırıcı kullanan birilerini kişisel olarak tanıyor musunuz?

- 01 ☐ Evet
02 ☐ Hayır

2.Kendiniz hiç yatıştırıcı kullandınız mı?

- 01 ☐ Evet
02 ☐☐☐ Hayır → Esrarla ilgili soruya gidin
03 ☐ Bilmiyorum → Esrarla ilgili soruya gidin

3.Son 12 ay boyunca, hiç yatıştırıcı kullandınız mı?

- 01 ☐ Evet
02 ☐ Hayır→ Esrarla ilgili soruya gidin

4.Son 30 gün boyunca, hiç yatıştırıcı kullandınız mı?

- 01 ☐ Evet
02 ☐ Hayır

5.Son 30 gün boyunca, kaç gün yatıştırıcı kullandınız ?

- 01 ☐ hergün veya hemen hergün
02 ☐ haftada birkaç kez
03 ☐ haftada en az bir kez
04 ☐ haftada bir kezden az

6.İlk defa yatıştırıcı ne zaman aldınız?

- 01 ☐ 15 yaşından önce
02 ☐ 15-20 yaş arası
03 ☐ 20-30 yaş arası
04 ☐ 30 yaşından sonra
05 ☐ Bilmiyorum, hatırlamıyorum

UÇUCU MADDE

1. Uçucu madde kullanan birilerini kişisel olarak tanıyor musunuz?

- 01 ☐ Evet
02 ☐ Hayır

2.Kendiniz hiç uçucu madde kullandınız mı?

- 01 ☐ Evet
02 ☐ ☐ ☐ Hayır → Esrarla ilgili soruya gidin
03 ☐ Bilmiyorum → Esrarla ilgili soruya gidin

3.Son 12 ay boyunca, hiç uęucu madde kullandınız mı?

- 01 ☐ Evet
02 ☐ Hayır→ Esrarla ilgili soruya gidin

4.Son 30 g¼n boyunca, hiç uęucu madde kullandınız mı?

- 01 ☐ Evet
02 ☐ Hayır

5.Son 30 g¼n boyunca, kaç g¼n uęucu madde kullandınız ?

- 01 ☐ herg¼n veya hemen herg¼n
02 ☐ haftada birkaç kez
03 ☐ haftada en az bir kez
04 ☐ haftada bir kezden az

6.İlk defa uęucu madde ne zaman aldınız?

- 01 ☐ 15 yaşımdan önce
02 ☐ 15-20 yaşı arası
03 ☐ 20-30 yaşı arası
04 ☐ 30 yaşımdan sonra
05 ☐ Bilmiyorum, hatırlamıyorum

ESRAR

1. Esrar kullanan birilerini kişisel olarak tanıyor musunuz?

- 01 ☐ Evet
02 ☐ Hayır

2.Kendiniz hiç esrar kullandınız mı?

- 01 ☐ Evet
02 ☐ ☐ ☐ Hayır → Esrarla ilgili soruya gidin
03 ☐ Bilmiyorum → Esrarla ilgili soruya gidin

3.Son 12 ay boyunca, hiç uç esrar kullandınız mı?

- 01 ☐ Evet
02 ☐ Hayır→ Esrarla ilgili soruya gidin

4.Son 30 gün boyunca, hiç esrar kullandınız mı?

- 01 ☐ Evet
02 ☐ Hayır

5.Son 30 gün boyunca, kaç gün esrarkullandınız ?

- 01 ☐ hergün veya hemen hergün
02 ☐ haftada birkaç kez
03 ☐ haftada en az bir kez
04 ☐ haftada bir kezden az

6.İlk defa esrar ne zaman aldınız?

- 01 ☐ 15 yaşından önce
02 ☐ 15-20 yaş arası
03 ☐ 20-30 yaş arası
04 ☐ 30 yaşından sonra
05 ☐ Bilmiyorum, hatırlamıyorum

BONZAI

1. Bonzai kullanan birilerini kişisel olarak tanıyor musunuz?

- 01 ☐ Evet
02 ☐ Hayır

2.Kendiniz hiç bonzai kullandınız mı?

- 01 ☐ Evet
02 ☐ ☐ ☐ Hayır → Esrarla ilgili soruya gidin
03 ☐ Bilmiyorum→ Esrarla ilgili soruya gidin

3.Son 12 ay boyunca, hiç uç bonzai kullandınız mı?

- 01 ☐ Evet

02 ☐ Hayır→ Esrarla ilgili soruya gidin

4.Son 30 gün boyunca, hiç bonzai kullandınız mı?

01 ☐ Evet

02 ☐ Hayır

5.Son 30 gün boyunca, kaç gün bonzaikullandınız ?

01 ☐ hergün veya hemen hergün

02 ☐ haftada birkaç kez

03 ☐ haftada en az bir kez

04 ☐ haftada bir kezden az

6.İlk defa bonzai ne zaman aldınız?

01 ☐ 15 yaşından önce

02 ☐ 15-20 yaş arası

03 ☐ 20-30 yaş arası

04 ☐ 30 yaşından sonra

05 ☐ Bilmiyorum, hatırlamıyorum

AMFETAMİNLER

1.Amfetaminler (speed,pep) kullanan birilerini kişisel olarak tanıyor musunuz?

01 ☐ Evet

02 ☐ Hayır

2. Kendiniz hiç amfetamin (speed,pep) kullandınız mı?

01 ☐ Evet

02 ☐ Hayır→ Ectasy'le ilgili soruya gidin

03 ☐ Bilmiyorum→ Ectasy'le ilgili soruya gidin

3.Son 12 ay boyunca, hiç amfetamin(speed,pep) kullandınız mı?

- 01 ☐ Evet
02 ☐ Hayır→ Ecstasy'le ilgili soruya gidin

4.Son 30 gün boyunca, hiç amfetamin (speed,pep) kullandınız mı?

- 01 ☐ Evet
02 ☐ Hayır→ Ecstasy'le ilgili soruya gidin

5. Son 30 gün boyunca, kaç gün amfetamin (speed, pep) kullandınız?

- 01 ☐ Her gün veya hemen her gün
02 ☐ Haftada birkaç kez
03 ☐ Haftada en az bir kez
04 ☐ Haftada bir kez den az

6.İlk defa amfetamin ne zaman aldınız?

- 01 ☐ 15 yaşından önce
02 ☐ 15-20 yaş arası
03 ☐ 20-30 yaş arası
04 ☐ 30 yaşından sonra
05 ☐ Bilmiyorum, hatırlamıyorum

ECTASY(XTC)

1. Ecstasy kullanan birilerini kişisel olarak tanıyor musunuz?

- 01 ☐ Evet

02 ☐ Hayır

2. Kendiniz hiç ecstasy kullandınız mı?

01 ☐ Evet

02 ☐ Hayır → Kokainle ilgili soruya gidin

03 ☐ Bilmiyorum → Kokainle ilgili soruya gidin

3. Son 12 ay boyunca, hiç ecstasy kullandınız mı?

01 ☐ Evet

02 ☐ Hayır → Kokainle ilgili soruya gidin

4. Son 30 gün boyunca, her ecstasy kullandınız mı?

01 ☐ Evet

02 ☐ Hayır → Kokainle ilgili soruya gidin

5. Son 30 gün boyunca, kaç gün ecstasy kullandınız?

01 ☐ Her gün veya hemen her gün

02 ☐ Haftada birkaç kez

03 ☐ Haftada en az bir kez

04 ☐ Haftada bir kez den az

6. İlk defa ecstasy ne zaman aldınız?

01 ☐ 15 yaşından önce

02 ☐ 15-20 yaş arası

03 ☐ 20-30 yaş arası

04 ☐ 30 yaşından sonra

05 ☐ Bilmiyorum, hatırlamıyorum

- 04 ☐ 30 yaşımdan sonra
05 ☐ Bilmiyorum, hatırlamıyorum

RELEVİN

1. Relevin kullanan birilerini kişisel olarak tanıyor musunuz?

- 01 ☐ Evet
02 ☐ Hayır

2. Kendiniz hiç relevin kullandınız mı?

- 01 ☐ Evet
02 ☐ Hayır→LSDyle ilgili soruya gidin
03 ☐ Bilmiyorum→ LSDyle ilgili soruya gidin

3. Son 12 ay boyunca, hiç relevin kullandınız mı?

- 01 ☐ Evet
02 ☐ Hayır→ LSDyle ilgili soruya gidin

4. Son 30 gün boyunca, hiç relevin kullandınız mı?

- 01 ☐ Evet
02 ☐ Hayır→ LSDyle ilgili soruya gidin

5. Son 30 gün boyunca, kaç gün relevin kullandınız?

- 01 ☐ Her gün veya hemen her gün
02 ☐ Haftada birkaç kez

03 ☐ Haftada en az bir kez

04 ☐ Haftada bir kez den az

6.İlk defa relevin ne zaman aldınız?

01 ☐ 15 yaşından önce

02 ☐ 15-20 yaş arası

03 ☐ 20-30 yaş arası

04 ☐ 30 yaşından sonra

05 ☐ Bilmiyorum, hatırlamıyorum

EROİN

1. Eroin kullanan birilerini kişisel olarak tanıyor musunuz?

01 ☐☐☐Evet

02 ☐ Hayır

2. Kendiniz hiç eroin kullandınız mı?

01 ☐ Evet

02 ☐ Hayır→Relevinle ilgili soruya gidin

03 ☐ Bilmiyorum→ Relevinle ilgili soruya gidin

3. Son 12 ay boyunca, hiç eroin kullandınız mı?

01 ☐ Evet

02 ☐ Hayır→ Relevinle ilgili soruya gidin

4. Son 30 gün boyunca, hiç eroin kullandınız mı?

01 ☐ Evet

02 ☐ Hayır→ Relewinle ilgili soruya gidin

5. Son 30 gün boyunca, kaç gün eroin kullandınız?

01 ☐ Her gün veya hemen her gün

02 ☐ Haftada birkaç kez

03 ☐ Haftada en az bir kez

04 ☐ Haftada bir kez den az

6. İlk defa eroin ne zaman aldınız?

01 ☐ 15 yaşından önce

02 ☐ 15-20 yaş arası

03 ☐ 20-30 yaş arası

04 ☐ 30 yaşından sonra

05 ☐ Bilmiyorum, hatırlamıyorum

LSD

1. LSD ('trips', 'asit') kullanan birilerini kişisel olarak tanıyor musunuz?

01 ☐ Evet

02 ☐ Hayır

2. Kendiniz hiç LSD ('trips', 'asit') kullandınız mı?

01 ☐ Evet

02 ☐ Hayır→Bonzai'yle ilgili soruya gidin

03 ☐ Bilmiyorum→ Bonzai'yle ilgili soruya gidin

3. Son 12 ay boyunca, hiç LSD ('trips', 'asit') kullandınız mı?

01 ☐ Evet

02 ☐ Hayır → Bonzai'yle ilgili soruya gidin

4. Son 30 gün boyunca, hiç LSD ('trips', 'asit') kullandınız mı?

01 ☐ Evet

02 ☐ Hayır → Bonzai'yle ilgili soruya gidin

5. Son 30 gün boyunca, kaç gün LSD ('trips', 'asit') kullandınız?

01 ☐ Her gün veya hemen her gün

02 ☐ Haftada birkaç kez

03 ☐ Haftada en az bir kez

04 ☐ Haftada bir kez den az

6. İlk defa LSD ('trips', 'asit') ne zaman aldınız?

01 ☐ 15 yaşından önce

02 ☐ 15-20 yaş arası

03 ☐ 20-30 yaş arası

04 ☐ 30 yaşından sonra

05 ☐ Bilmiyorum, hatırlamıyorum

KODEİNLİ ŞURUP

1. Kodeinli şurup kullanan birilerini kişisel olarak tanıyor musunuz?

- 01 ☐ Evet
- 02 ☐ Hayır

2. Kendiniz hiç kodeinli şurup kullandınız mı?

- 01 ☐ Evet
- 02 ☐ ☐ ☐ Hayır → Esrarla ilgili soruya gidin
- 03 ☐ Bilmiyorum → Esrarla ilgili soruya gidin

3. Son 12 ay boyunca, hiç kodeinli şurup kullandınız mı?

- 01 ☐ Evet
- 02 ☐ Hayır → Esrarla ilgili soruya gidin

4. Son 30 gün boyunca, hiç kodeinli şurup kullandınız mı?

- 01 ☐ Evet
- 02 ☐ Hayır

5. Son 30 gün boyunca, kaç gün kodeinli şurup kullandınız ?

- 01 ☐ hergün veya hemen hergün
- 02 ☐ haftada birkaç kez
- 03 ☐ haftada en az bir kez
- 04 ☐ haftada bir kezden az

6. İlk defa kodeinli şurup ne zaman aldınız?

- 01 ☐ 15 yaşından önce
- 02 ☐ 15-20 yaş arası
- 03 ☐ 20-30 yaş arası
- 04 ☐ 30 yaşından sonra
- 05 ☐ bilmiyorum, hatırlamıyorum

Hap kullanımı

1. Hap (akineton, roş-rohypnol, nembotal-sarı bomb vs.) kullanan birilerini kişisel olarak tanıyor musunuz?

- 01 ☐ Evet
- 02 ☐ Hayır

2. Kendiniz hiç hap (akineton, roş-rohypnol, nembotal-sarı bomb vs.) kullandınız mı?

- 01 ☐ Evet
- 02 ☐ Hayır→Bonzai'yle ilgili soruya gidin
- 03 ☐ Bilmiyorum→ Bonzai'yle ilgili soruya gidin

3. Son 12 ay boyunca, hiç hap (akineton, roş-rohypnol, nembotal-sarı bomb vs.) kullandınız mı?

- 01 ☐ Evet
- 02 ☐ Hayır → Bonzai'yle ilgili soruya gidin

4. Son 30 gün boyunca, hiç hap (akineton, roş-rohypnol, nembotal-sarı bomb vs.) kullandınız mı?

01 ☐ Evet

02 ☐ Hayır → Bonzai'yle ilgili soruya gidin

5. Son 30 gün boyunca, kaç gün hap (akineton, roş-rohypnol, nembotal-sarı bomb vs.) kullandınız?

01 ☐ Her gün veya hemen her gün

02 ☐ Haftada birkaç kez

03 ☐ Haftada en az bir kez

04 ☐ Haftada bir kez den az

6. İlk defa hap (akineton, roş-rohypnol, nembotal-sarı bomb vs.) ne zaman aldınız?

01 ☐ 15 yaşından önce

02 ☐ 15-20 yaş arası

03 ☐ 20-30 yaş arası

04 ☐ 30 yaşından sonra

05 ☐ Bilmiyorum, hatırlamıyorum

Anabolizan Steroid

1. Anabolizan steroid kullanan birilerini kişisel olarak tanıyor musunuz?

01 ☐ Evet

02 ☐ Hayır

2. Kendiniz hiç anabolizan steroid kullandınız mı?

- 01 ☐ Evet
- 02 ☐ Hayır→Relevanle ilgili soruya gidin
- 03 ☐ Bilmiyorum→ Relevanle ilgili soruya gidin

3. Son 12 ay boyunca, hiç anabolizan steroid kullandınız mı?

- 01 ☐ Evet
- 02 ☐ Hayır→ Relevanle ilgili soruya gidin

4. Son 30 gün boyunca, hiç anabolizan steroid kullandınız mı?

- 01 ☐ Evet
- 02 ☐ Hayır→ Relevanle ilgili soruya gidin

5. Son 30 gün boyunca, kaç gün eroin kullandınız?

- 01 ☐ Her gün veya hemen her gün
- 02 ☐ Haftada birkaç kez
- 03 ☐ Haftada en az bir kez
- 04 ☐ Haftada bir kez den az

6.İlk defa anabolizan steroid ne zaman aldınız?

- 01 ☐ 15 yaşından önce
- 02 ☐ 15-20 yaş arası
- 03 ☐ 20-30 yaş arası
- 04 ☐ 30 yaşından sonra
- 05 ☐ Bilmiyorum, hatırlamıyorum

CURRICULUM VITAE

1. Personal Information:

NAME, SURNAME:	ELİF CEREN SERT
BIRTH DATE AND PLACE:	20.06.1990 , İstanbul
Duty:	Psychologist
Telephone:	05338382685
e-mail:	elifcerensert@gmail.com

2. Education

Level	Field	University	Year
Bachelor	Psychology	Near East University	2013

3. Professional Experience

Period of Duty	Title	Field	Work Place
2014-2015	Intern Clinical Psychologist	Clinical Psychology	Bakırköy Ruh ve Sinir Hastalıkları Hastahanesi
2014-2015	Intern Clinical Psychologist	Clinical Psychology	Barış Ruh ve Sinir Hastalıkları Hastahanesi