



**CONFLUENT HYPERGEOMETRIC FUNCTION
WITH KUMMER'S FIRST FORMULA**

**A THESIS SUBMITTED TO THE GRADUATE
SCHOOL OF APPLIED SCIENCES
OF
NEAR EAST UNIVERSITY**

**By
SHWAN SWARA FATAH**

**In Partial Fulfillment of the Requirements for
the Degree of Master of Science
in
Mathematics**

NICOSIA, 2016

**Shwan Swara Fatah: CONFLUENT HYPERGEOMETRIC FUNCTION WITH
KUMMER'S FIRST FORMULA**



**Approval of Director of Graduate School of
Applied Sciences**

A blue ink signature of Prof. Dr. İlkey SALİHOĞLU, written over a circular blue stamp that reads "NEAR EAST UNIVERSITY" and "APPLIED SCIENCES".

Prof. Dr. İlkey SALİHOĞLU

**We certify that, this thesis is satisfactory for the award of the degree of Master of
Sciences in Mathematics.**

Examining Committee in Charge:

A blue ink signature of Prof. Dr. Adigüzel Dosiyevev.

Prof. Dr. Adigüzel Dosiyevev

**Committee Chairman,
Department of Mathematics,
Near East University**

A blue ink signature of Assist. Prof. Burak Şekeroğlu.

Assist. Prof. Burak Şekeroğlu

**Supervisor,
Department Mathematics,
Near East University.**

A blue ink signature of Dr. Emine Çeliker.

Dr. Emine Çeliker

**Mathematics Research and Teaching
Group Middle East Technical
University, Northern Cyprus Campus**

I hereby declare that all information in this document has been obtained and presented in accordance with academic rules and ethical conduct. I also declare that, as required by these rules and conduct, I have fully cited and referenced all material and results that are not original to this work.

Name, Last name: Shwan S Ward Patah

Signature: 

Date: 21/4/2016

ACKNOWLEDGMENTS

I wish to express my profound gratitude to my supervisor Assist. Prof. Dr. Burak Şekeroğlu for his time, and corrections which have contributed vastly to the completion of this work. I owe a lot of gratitude to the Chairman of the Committee Assoc. Prof. Dr. Evren Hınçal for his fatherly advice in making this work a success.

I would also like to thank Dr. Abdurrahman for his corrections and advice towards making my research a success.

I wish to also express my gratitude to all staff of Mathematics Department of Near East University for their advice, support and the vast knowledge I have acquired from them. Their excitement and willingness to provide feedback made the completion of this research an enjoyable experience.

I am indeed most grateful to my late father, Mr Swara, my mother Mrs Shamsa whose constant prayers, love, support and guidance have been my source of strength and inspiration throughout my years of study.

I cannot forget to acknowledge the support I received from my beloved siblings, especially brother Muhammad and his wife Shawnm, sister Hasiba who stood by me throughout the stormy years and gave me the courage that I very much needed to pursue my studies.

I also wish to acknowledge all my friends especially Musa Dan-azumi Mohammed who helped me through with this research and relatives whose names are too numerous to mention.

To my parents...

ABSTRACT

The study is an examination of the definitions and basic properties of hypergeometric function, confluent hypergeometric function, The main objective of the study confluent hypergeometric function with Kummer's first formula. Several properties such as contiguous function relations., differential equations and Elementary series manipulation for these hypergeometric and confluent hypergeometric families are obtained. Was to ascertain an approximation of solution of confluent hypergeometric function. Were therefore drawn from the study that the Kummer's function has wide application in various subjects and hence proving stability or other properties were also drawn to be of paramount importance.

Keywords: Hypergeometric function; Confluent hypergeometric function; Kummer's first formula; Gamma function; Pochhammer function

ÖZET

çalışma, Hipergeometrik fonksiyonların ve Birleşik hyperbolic fonksiyonların tanımlarını ve temel özelliklerini inceler. Çalışmanın temel amacı birinci Kummer formula ile birleşik hipergeometrik fonksiyonları çalışmaktır. Hipergeometrik ve birleşik hipergeometrik ailelerinin differensiyel denklemleri, bitişik fonksiyon ilişkileri, ve temel seri manipulasyonları gibi bazı özellikler elde edilmiştir. Birleşik hipergeometrik fonksiyon çözümlerinin yaklaşımları bulunmuştur. Kummer's fonksiyonları çeşitli konlarda geniş uygulama alanlarına sahiptir ve kararlılığı ve diğer özelliklerinin de oldukça önemli olduğu bu tezde vurgulanmıştır.

Anahtar Kelimeler: Hipergeometrik fonksiyon; Birleşik Hipergeometrik fonksiyon; Kummer in birinci formula; Gamma fonksiyon; Pochhammer fonksiyon

TABLE OF CONTENTS

ACKNOWLEDGMENTS.....	i
ABSTRACT	iii
ÖZET	iv
TABLE OF CONTENTS	v
LIST OF SYMBOLS	viii

CHAPTER 1: INTRODUCTION

1.2 Gamma Function	2
1.2.1 Definition	2
1.2.2 Some basic properties of Gamma function with their proofs	2
1.2.3 Lemma	3
1.3 Definition.....	3
1.3.1 Theorem	3
1.3.2 Some special values for Beta function.....	4
1.4 Definition	4
1.4.1 Some properties of Pochhammer symbol	4
1.4.2 Theorem	5
1.4.3 Lemma	6

CHAPTER 2: CONFLUENT HYPERGEOMETRIC FUNCTION

2.1 Hypergeometric Function	7
2.1.1 Definition	7
2.1.2 Functions with representations like Hypergeometric series	8
2.1.3 Properties of Hypergeometric functions	8
2.1.3.1 Differential representation.....	8
2.1.3.2 Integral representation	8
2.1.3.3 The Hypergeometric equation	9
2.1.4 Problem	10
2.2. Generalized Hypergeometric Function.....	10
2.3 Bessel Function	12
2.3.1 Definition	12

2.4 Confluent Hypergeometric Function	12
2.4.1 Definition	12
2.4.2 Relation to other functions	13
2.4.3 Theorem	13
2.4.4 Elementary properties of Confluent Hypergeometric Function.....	14
2.4.4.1 Differential representation.....	14
2.4.4.2 Integral representation	14
2.4.4.3 Theorem.....	15
2.4.4.5 Confluent Hypergeometric equation	15
2.4.4.6 Multiplication formula	16
2.4.4.7 The Contiguous function relation.....	16
2.4.5 Some example of confluent Hypergeometric function	17

CHAPTER 3: CONFLUENT HYPERGEOMETRIC FUNCTION WITH KUMMER FIRST FORMULA

3.1 Theorem	20
3.2 Theorem	21
3.3 Some example of Kummer's formula with (CHF)	23

CHAPTER 4: SEVERAL PROPERTIES OF HYPERGEOMETRIC FUNCTION

4.1 Properties	25
4.1.1 The Contiguous Function relations.....	25
4.1.2 Hypergeometric differential equation:.....	27
4.1.3 Elementary series manipulation	29
4.1.5 A quadratic transformation	30
4.1.5.1 Theorem.....	30
4.1.6 Additional properties	33
4.1.6.1 Theorem.....	33
4.2 Some Theorem Without Proof.....	34
4.2.1 Theorem	34
4.2.2 Theorem	34
4.2.3 Theorem	34

CHAPTER 5: CONCLUSION AND SUGGESTIONS FOR FUTURE STUDIES..35

REFERENCES 36

LIST OF SYMBOLS

$\Gamma(x)$	Gamma Functions
$(x)_n$	Pohammer Symbol
${}_2F_1(a, b; c; z)$	Hypergeomtric function
${}_pF_q\left(\begin{matrix} a_1, a_2, a_3, \dots, a_p \\ b_1, b_2, b_3, \dots, b_q \end{matrix}; z\right)$	General hypergeomtric function
${}_1F_1(a; c; z)$	Confluent hypergeomtric function
${}_0F_0(-, z)$	Expositional function
${}_1F_0\left(\begin{matrix} a \\ - \end{matrix}, z\right)$	BinoType equation here.mial function
J_α	Bessel function of the first kind
\prod	Product
$[\]$	Bracket

CHAPTER 1

INTRODUCTION

This chapter outlines several basic definitions, theorems and some properties of special functions. This study thrives to proffer insights about the confluent hypergeometric function with by employing the Kummer's formula. The notion behind the Kummer confluent hypergeometric function (CHF) stems from an essential category of special functions of mathematical physics. Kummer's formula in (CHF) can be decomposed into the following;

The initial Kummer's formula assumes the following form:

$$e^{-z} {}_1F_1(a; c; z) = \sum_{n=0}^{\infty} \frac{(c-a)_n (-z)^n}{(c)_n n!} = {}_1F_1(c-a; c; -z), c \neq \{0\} \cup \{-1, -2, -3, \dots\}$$

And, kummer's second formula

$$e^{-z} {}_1F_1(a, 2a; 2z) = \sum_{n=0}^{\infty} \frac{(\frac{1}{4}z^2)^n}{(a + \frac{1}{2})_n n!} = {}_0F_1\left(-; a + \frac{1}{2}; \frac{1}{4}z^2\right).$$

if a is not odd positive integer.

This study will therefore offer further explanations about the Kummer's first formula in confluent hyper geometric functions. Despite the fact that Gauss played an essential role in the systematic study of the hypergeometric function, (Kummer, 1837) assumed a critical role in the development of properties of confluent hypergeometric functions. Kummer published his work on this function in 1836 and since that time it has been commonly referred to as the Kummer's function (Andrews, 1998). Under the hypergeometric function, the confluent hypergeometric function is related to a countless number of different functions.

This work therefore outlines the general and basic properties of hypergeometric and confluent hypergeometric function and the Kummer's first formula. This study will also extend to incorporate the related examples and theorems.

The first chapter deals with the synopsis of basic definitions, theorems and exceptions of the hyper geometric functions while the second chapter is a blueprint of definitions, properties and theorems of confluent hyper geometric functions. Meanwhile, chapter three lays out examples and special cases of Kummer's first formula coupled with reinforcing explanations. A recapitulation of properties of the hypergeometric functions is given in the fourth chapter while the fifth chapter concludes this study by looking at conclusions that can be drawn from this study.

1.2 Gamma Function

It is undoubtable that most essential functions in applied sciences are defined via improper integrals. Of notable effect is Gamma functions. Such functions have several applications in Mathematics and Mathematical Physics.

1.2.1 Definition

The elementary definition of the gamma function is Euler's integral (Gogolin, 2013)

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt.$$

is converges for any $z > 0$

1.2.2 Some basic properties of Gamma function with their proofs(Özergin, 2011).

$$\Gamma(1) = \int_0^{\infty} e^{-t} dt = -e^{-t} \Big|_0^{\infty} = 1$$

$$\Gamma\left(\frac{1}{2}\right) = \int_0^{\infty} \frac{e^{-t}}{\sqrt{t}} dt = 2\Gamma(1) = 2 \int_0^{\infty} e^{-u^2} du = 2 \frac{\sqrt{\pi}}{2} = \sqrt{\pi}$$

$$\Gamma(x+1) = \int_0^{\infty} t^x e^{-t} dt = -t^x e^{-t} \Big|_0^{\infty} - \int_0^{\infty} x t^{x-1} (-e^{-t}) dt = x \int_0^{\infty} t^{x-1} e^{-t} dt = x \Gamma(x)$$

1.2.3 Lemma

The Gamma function satisfies the functional equation

$$\Gamma(x+1) = x \Gamma(x), \quad x > 0$$

Moreover, by iteration for $x > 0$ and $n \in \mathbb{N}$

$$\Gamma(x+n) = \Gamma(x+n-1) \cdot \dots \cdot (x+1)x \Gamma(x) = \prod_{i=1}^n (x+1-i) \Gamma(x)$$

$$\Gamma(n+1) = \left(\prod_{i=1}^n i \right) \Gamma(1) = \prod_{i=1}^n (i) = n!$$

In other words, the Gamma function can be interpreted as an extension of factorials.

1.3 Definition (Sebah, 2002)

The beta function or Eulerian integral of the first kind is given by

$$B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt, \quad \text{where } x, y > 0$$

This definition is also valid for complex numbers x and y such as

$$R(x) > 0 \text{ and } R(y) > 0$$

1.3.1 Theorem (Gronan, 2003)

$$\text{if } R(x) > 0 \text{ and } R(y) > 0 \text{ then } B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} = B(y, x)$$

1.3.2 Some special values for Beta function

$$B\left(\frac{1}{2}, \frac{1}{2}\right) = \pi ,$$

$$B(x, 1) = \frac{1}{x} ,$$

$$B(x, n) = \frac{(n-1)!}{x(x+1)\dots(x+n-1)} \quad n \geq 1$$

1.4 Definition

Let x be a real or complex number and n be a positive integer,

$$(x)_n = \frac{\Gamma(x+n)}{\Gamma(x)} = x(x+1)\dots(x+n-1)$$

“Pochhammer Symbol” is where $(x)_n$ is used to represent the falling factorial sometimes called the descending factorial, falling sequential product, lower factorial (Freedman, 2013).

1.4.1 Some properties of Pochhammer symbol

$$i) (a+n)_k = \frac{(a)_{n+k}}{(a)_n},$$

Where a is a real or complex number and n, k are natural numbers

$$ii) \frac{(a)_{2k}}{2^{2k}} = \left(\frac{a}{2}\right)_k \left(\frac{a}{2} + \frac{1}{2}\right)_k$$

Where a is a complex number and k is a natural number

$$iii) \frac{(2k)!}{2^{2k}k!} = \left(\frac{1}{2}\right)_k$$

Where k : is a natural number.

Note

If $a = 1$ then we have $(a)_n = (1)_n = 1 \times 2 \times 3 \times \dots \times n = n!$

If $a = 2$ then $(2)_n = (n+1)$ and also we have

$$(a)_n = (-N)_n = (-N)(-N+1)(-N+2) \cdots (-N+n-1) = 0 \text{ if } a = -n, \\ n = \{0, 1, 2, \dots\}.$$

$$(a)_0 = 1, \quad a \neq 0$$

1.4.2 Theorem

Show that for $0 \leq k \leq n$,

$$(a)_{n-k} = \frac{(-1)^k (a)_n}{(1-a-n)_k}$$

Note particularly the special case $a = 1$

Proof:

Consider $(a)_{n-k}$ for $0 \leq k \leq n$,

$$\begin{aligned} (a)_{n-k} &= a(a+1) \dots (a+n-k-1) \\ &= \frac{a(a+1) \dots (a+n-k-1)[(a+n-k)(a+n-k+1) \dots (a+n-1)]}{(a+n-k)(a+n-2) \dots (a+n-k)} \\ &= \frac{(a)_n}{(a+n-k)_k} \\ &= \frac{(a)_n}{(-1)^k (1-a-n)_k} \\ &= \frac{(-1)^k (a)_n}{(1-a-n)_k} \end{aligned}$$

$$\text{Not for } a = 1, (n-k)! = \frac{(-1)^k n!}{(-n)_k}.$$

1.4.3 Lemma

$$(\alpha)_{2n} = 2^{2n} \left(\frac{\alpha}{2}\right)_n \left(\frac{\alpha+1}{2}\right)_n$$

Proof:

$$\begin{aligned} (\alpha)_{2n} &= \alpha(\alpha+1)(\alpha+2) \\ &= 2^{2n} \left(\frac{\alpha}{2}\right) \left(\frac{\alpha+1}{2}\right) \left(\frac{\alpha}{2}+1\right) \dots \left(\frac{\alpha}{2}+n-1\right) \left(\frac{\alpha+1}{2}+n-1\right) \\ &= 2^{2n} \left(\frac{\alpha}{2}\right) \left(\frac{\alpha+1}{2}\right) \left(\frac{\alpha}{2}+1\right) \left(\frac{\alpha}{2}+n-1\right) \left(\frac{\alpha+1}{2}\right) \left(\frac{\alpha+1}{2}+1\right) \dots \left(\frac{\alpha+1}{2}+n-1\right) \\ &= 2^{2n} \left(\frac{\alpha}{2}\right)_n \left(\frac{\alpha+1}{2}\right)_n \end{aligned}$$

CHAPTER 2

CONFLUENT HYPERGEOMETRIC FUNCTION

This section draws attention on the confluent hypergeometric function, its definition and inherent properties. Due to the importance that is attached to the confluent hypergeometric function in hypergeometric function; this study will therefore draw attention to the examination of the hyper geometric function.

2.1 Hypergeometric Function

The function ${}_2F_1(a, b; c; x)$ corresponding to $p=2, q=1$ is the first hyper geometric function to be examined (and, in general, emerges in prominence especially in physical problems), as is synonymously referred to as "the" hyper geometric equation or, more explicitly, Gauss's hyper geometric function (Gauss, 1812; Barnes 1908). To confound matters much more, the term "hyper geometric function" is less usually used to mean shut structure, and "hyper geometric series" is sometimes used to mean hyper geometric function.

Hyper geometric functions are solutions to the hyper geometric differential equation, which has a regular singular point at the starting point. A hyper geometric function can be derived from the hyper geometric differential equation.

2.1.1 Definition

(Rainville, 1965). Asserts that a hyper geometric function can be defined as follows;

$$F(a, b; c; z) = {}_2F_1(a, b; c; z) = F(b, a; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n n!} z^n, |z| \leq 1 \quad (2.1)$$

For c neither zero nor negative integer. In 2.1, the notation

1 - Refers to number of parameters in denominator

2 - Refers to number of parameters in numerator

2.1.2 Functions with representations like Hypergeometric series

$$F(1, b, b, z) = \sum_{n=0}^{\infty} \frac{(1)_n}{n!} \cdot z^n = \sum_{n=0}^{\infty} z^n$$

$$\arcsin z = F\left(\frac{1}{2}, \frac{1}{2}; \frac{3}{2}; z^2\right)$$

$$\ln(1+z) = \sum_{n=0}^{\infty} \frac{(-1)^n}{n+1} z^n = \sum_{n=0}^{\infty} \frac{(1)_n (1)_n}{(2)_n} \frac{(-1)^n z^{n+1}}{n!} = zF\left(\frac{1}{2}, \frac{1}{2}; -z\right)$$

2.1.3 Properties of Hypergeometric functions

2.1.3.1 Differential representation

The Differential representation of the hypergeometric function is given by

$$\begin{aligned} \frac{d}{dz} F(a, b; c; z) &= \sum_{n=1}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{z^{n-1}}{(n-1)!} \\ &= \sum_{n=0}^{\infty} \frac{(a)_{n+1} (b)_{n+1} z^n}{(c)_{n+1} n!} \\ &= \frac{ab}{c} \sum_{n=0}^{\infty} \frac{(a+1)_n (b+1)_n z^n}{(c+1)_n n!} \\ &= \frac{ab}{c} F(a+1, b+1; c+1; z) \end{aligned}$$

2.1.3.2 Integral representation

$$F(a, b; c; z) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 t^{b-1} (1-t)^{c-b-1} (1-xt)^a dt \quad c > b > 0$$

Where Gamma is defined by

$$\Gamma(x) = \int_0^{\infty} t^x e^{-t} dt, x > 0$$

2.1.3.3 The Hypergeometric equation

The linear second-order DE

$$z(1-z) \frac{dw^2}{dz^2} + (c - (\alpha + b + 1)z) \frac{dw}{dz} - abw = 0$$

is called the hypergeometric equation

These functions were studied by numerous mathematicians including Riemann who gathered in their conduct as functions of a complex variable, also, concentrated on its analytic continuation regarding it as a solution to the differential equation (Campos, 2001).

$$z(1-z) \frac{dw^2}{dz^2} + (c - (\alpha + b + 1)z) \frac{dw}{dz} - abw = 0 \quad (2.1)$$

or, multiplying equation (4) by z and denoting $\theta = z \frac{d}{dz}$,

$$[\theta(\theta + c - 1) - z(\theta + \alpha)(\theta + b)](z) = 0 \quad (2.2)$$

Equation 2.1 or 2.2, has three regular singular points at 0, 1 and ∞ , and it is

Up to standardization the general form of a second order linear differential equation with this conduct.

Note if one of the numerator parameters a or b are equal to the denominator parameter c we get

$$\begin{aligned} {}_2F_1 \left(\begin{matrix} a, b \\ b \end{matrix} ; z \right) &= \sum_{n=0}^{\infty} \frac{(b)_n (a)_n}{(b)_n n!} z^n \\ &= {}_1F_0 \left(\begin{matrix} a \\ - \end{matrix} ; z \right) \\ &= \sum_{n=0}^{\infty} \binom{-a}{n} (-z)^n \\ &= (1-z)^{-a}, \quad |z| < 1 \end{aligned}$$

2.1.4 Problem

Which results in

$$F \left[\begin{matrix} -n, b; \\ c; \end{matrix} 1 \right] = \frac{(c-b)_n}{(c)_n}$$

Solution

Consider $F(-n, b; c; 1)$. at once, if $R(c-b) > 0$,

$$F(-n, b; c; 1) = \frac{\Gamma(c)\Gamma(c-b+n)}{\Gamma(c+n)\Gamma(c-b)} = \frac{(c-b)_n}{(c)_n}$$

Actually the condition $R(c-b) > 0$ is not necessary because of the termination of the series involved.

2.2. Generalized Hypergeometric Function

As outlined in the definition (1) there are two numerator parameters, a and b ; and one denominator, c . it is a natural generalization to move from the definition (1) to a similar function with any number of numerator and denominator parameters.

We define a generalized hyper geometric function by

$$\begin{aligned} {}_pF_q \left(\begin{matrix} a_1, a_2, a_3, \dots, a_p; \\ b_1, b_2, b_3, \dots, b_q; \end{matrix} z \right) &= \sum_{n=0}^{\infty} \frac{(a_1)_n (a_2)_n (a_3)_n \dots (a_p)_n z^n}{(b_1)_n (b_2)_n (b_3)_n \dots (b_q)_n n!} \\ &= 1 + \sum_{n=1}^{\infty} \frac{\prod_{i=1}^p (a_i)_n z^n}{\prod_{i=1}^q (b_i)_n n!} \quad i = 1, 2, 3, \dots, n \end{aligned}$$

The parameters must be such that the denominator factors in the terms of the series are never zero. When one of the numerator parameters a_i equals $-N$, where N is a

nonnegative integer, the hypergeometric function is a polynomial in z (see below). Otherwise, the radius of convergence p of the hypergeometric series is given by

$$p = \begin{cases} \infty & \text{if } p < q + 1 \\ 0 & \text{if } p > q + 1 \\ 1 & \text{if } p = q + 1 \end{cases}$$

This follows directly from the ratio test. In fact, we have

$$\lim_{n \rightarrow \infty} \left| \frac{c_{n+1}}{c_n} \right| = \begin{cases} \infty & \text{if } p < q + 1 \\ 0 & \text{if } p > q + 1 \\ 1 & \text{if } p = q + 1 \end{cases}$$

In the case that $p = q + 1$ the situation that $|z| = 1$ is of special interest.

The hypergeometric series ${}_qF_p (a_1, a_2, \dots, a_{q+1}; b_1, b_2, \dots, b_p, z)$

with $|z| = 1$ converges absolutely if $\operatorname{Re} (\sum b_i - \sum a_j) \leq 0$

The series converges conditionally if $|z| = 1$ with $z \neq 1$ and $-1 < \operatorname{Re} (\sum b_i - \sum a_j) \leq 0$

And the series diverges if $\operatorname{Re} (\sum b_i - \sum a_j) \leq -1$.

Two elementary instances of the ${}_pF_q$ follow if no numerator or denominator parameters are present.

Which results to

$${}_0F_0 \left(\begin{matrix} - \\ - \end{matrix} ; z \right) = \sum_{n=0}^{\infty} \frac{z^n}{n!} = e^z$$

Which is called the exponential function where $z \in \mathbb{C}$

And also if we have one numerator parameter without denominator parameter, we obtain

$${}_1F_0 \left(\begin{matrix} a \\ - \end{matrix} ; z \right) = \sum_{n=0}^{\infty} \frac{(a)_n z^n}{n!} = (1 - z)^{-a}, \quad z \in \mathbb{C}$$

is called a binomial function

2.3 Bessel Function

We already know that the ${}_0F_0$ is an exponential and that ${}_1F_0$ is a binomial. It is natural to examine next the most general ${}_0F_1$, the only other ${}_pF_q$ with less than two parameters. The function we shall study is not precisely the ${}_0F_1$ but one that has an extra factor definition below (Dickenstein, 2004).

2.3.1 Definition

If n is not a negative integer

$$J_n(z) = \frac{\left(\frac{z}{2}\right)^n}{\Gamma(1+n)} {}_1F_0\left(-; 1+n; -\frac{z^2}{4}\right).$$

2.4 Confluent Hypergeometric Function

This section provides an examination of the most powerful methods implemented to accurately and efficiently evaluate the confluent hypergeometric function, Kummer's (confluent hypergeometric) function $M(a, b, z)$, introduced by (Kummer, 1837),

The term confluent refers to the merging of singular points of families of differential equations; confluent is Latin for "to flow together."

2.4.1 Definition

The Kummer confluent hypergeometric function is defined by the absolutely convergent infinite power series"

$$M(a, c, z) = {}_1F_1\left(\begin{matrix} a \\ c \end{matrix}; z\right) = \sum_{n=0}^{\infty} \frac{(a)_n z^n}{(c)_n n!}, \quad -\infty < z < \infty$$

It is analytic, regular at zero entire single-valued transcendental function of all a, c, x , (real or complex) except $c \neq 0$ or a negative integer.

Note

The confluent hypergeometric function is related to the hypergeometric function according to

$$\lim_{b \rightarrow \infty} \left(\begin{matrix} a, b \\ c \end{matrix} ; \frac{z}{b} \right) = \lim_{b \rightarrow \infty} \sum_{n=0}^{\infty} \frac{(a)_n (b)_n \left(\frac{z}{b}\right)^n}{(c)_n n!} = \sum_{n=0}^{\infty} \frac{(a)_n z^n}{(c)_n n!} \lim_{b \rightarrow \infty} \frac{(b)_n}{b^n} = \sum_{n=0}^{\infty} \frac{(a)_n z^n}{(c)_n n!}$$

So that $\lim_{b \rightarrow \infty} \left(\begin{matrix} a, b \\ c \end{matrix} ; \frac{z}{b} \right) = m(a, c, z)$

2.4.2 Relation to other functions

i) $m(-n; 1; z) = \ln(z)$

ii) $m(a; a, z) = e^z$

iii) $m(1, ; 2; 2z) = \frac{e^z}{z} \sinh z$

2.4.3 Theorem

$$m(a; a, z) = e^z$$

Proof:

$$m(a, a, z) = \sum_{n=0}^{\infty} \frac{(a)_n z^n}{(a)_n n!} = \sum_{n=0}^{\infty} \frac{z^n}{n!} = e^z$$

2.4.4 Elementary properties of Confluent Hypergeometric Function

2.4.4.1 Differential representation

Because of the similarity of definition to that of $F(a, b; c; z)$, the function $M(a; c; z)$ obviously has many properties analogous to those of the hypergeometric function (ko, 2011).

For example, it is easy to show that;

$$i) \frac{d}{dz} m(a, c; z) = \frac{a}{c} m(a+1, c+1; z).$$

Since

$$\begin{aligned} \frac{d}{dz} m(a, c; z) &= \sum_{n=1}^{\infty} \frac{(a)_n}{(c)_n} \frac{z^{n-1}}{(n-1)!} = \sum_{n=0}^{\infty} \frac{(a)_{n+1} z^n}{(c)_{n+1} n!} \\ &= \frac{a}{c} \sum_{n=0}^{\infty} \frac{(a+1)_n z^n}{(c+1)_n n!} \\ &= \frac{a}{c} m(a+1; c+1; z). \end{aligned}$$

Also in general

$$ii) \frac{d^k}{dz^k} m(a, c; z) = \frac{(a)_k}{(c)_k} m(a+k, c+k; z), \quad k = 1, 2, 3, \dots$$

2.4.4.2 Integral representation

Based on Euler's integral representation for the ${}_2F_1$ hypergeometric function, one might expect that the confluent hypergeometric function satisfies

$$\begin{aligned} m(a; c; z) &= {}_1F_1 \left(\begin{matrix} a \\ c \end{matrix} ; z \right) \\ &= \lim_{b \rightarrow \infty} \left(\begin{matrix} a, b \\ c \end{matrix} ; \frac{z}{b} \right) \end{aligned}$$

$$= \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^1 e^{zt} t^{a-1} (1-t)^{c-a-1} dt, \quad c > a > 0$$

2.4.4.3 Theorem

For $\text{Re } c > \text{Re } a > 0$ we have

$${}_1F_1\left(\begin{matrix} a \\ c \end{matrix}; z\right) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^1 e^{zt} t^{a-1} (1-t)^{c-a-1} dt$$

Proof: note that we have

$$\int_0^1 e^{xt} t^{a-1} (1-t)^{c-a-1} dt = \sum_{n=1}^{\infty} \frac{z^n}{n!} \int_0^1 t^{n+a-1} (1-t)^{c-a-1} dt$$

And $\text{Re } a > 0$

$$e(c-a) > 0$$

$$\int_0^1 t^{n+a-1} (1-t)^{c-a-1} dt = B(n+a, c-a)$$

$$= \frac{\Gamma(n+a)\Gamma(c-a)}{\Gamma(n+c)}$$

$$= \frac{\Gamma(a)\Gamma(c-a)}{\Gamma(c)} \frac{(a)_n}{(c)_n}$$

for $n = 0, 1, 2, \dots$, this implies that

$$\frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^1 e^{zt} t^{a-1} (1-t)^{c-a-1} dt = \sum_{n=0}^{\infty} \frac{(a)_n z^n}{(c)_n n!} = {}_1F_1\left(\begin{matrix} a \\ c \end{matrix}; z\right).$$

2.4.4.5 Confluent Hypergeometric equation

The Confluent hypergeometric equation established by (Buchholz, 2013) defines the hypergeometric function $y = F(a, b; c; z)$ as a solution of Gauss' equation

$$z(1-z) \frac{d^2 w}{dz^2} + (c - (\alpha + b + 1)z) \frac{dw}{dz} - \alpha b w = 0 \quad (2.3)$$

By making the change of variable $z = \frac{x}{b}$ (2.6) becomes

$$\left(1 - \frac{x}{b}\right)w'' + \left(c - x - \frac{a+1}{b}x\right)w' - aw = 0$$

and then allowing $b \rightarrow \infty$ we find

$$xw'' + (c - x)w' - aw = 0 \quad (2.4)$$

For $C \notin \mathbb{Z}$ the general solution of the confluent hypergeometric differential equation (2.4) can be written as

$$w(z) = A {}_1F_1\left(\frac{a}{c}; z\right) + Bz^{1-c} {}_1F_1\left(\frac{a+1-c}{2-c}; z\right)$$

with A and B arbitrary constants

2.4.4.6 Multiplication formula

A known formula, given by (Luke, 2014) can be utilized to determine the value of the confluent hypergeometric function in terms of another confluent hypergeometric function with the same parameters but with the variable of opposite sign. This formula can be specified as follows;

$${}_1F_1(a; b; z) \times {}_1F_1(a; b; -z) = {}_2F_3(a, b - a; b, \frac{1}{2}b + \frac{1}{2}; \frac{z^2}{4})$$

2.4.4.7 The Contiguous function relation

The function $m(a; c; z)$ also satisfies recurrence relations involving the contiguous functions $m(a \pm 1; c; z)$ and $m(a; c \pm 1; z)$. from these four contiguous functions, taken two at a time, we find six recurrence relations with coefficients at most linear in z (Pearson, 2009).

- i) $(c - a - 1)m(a; c; z) + am(a + 1; c; z) = (c - 1)m(a; c - 1; z)$
- ii) $cm(a; c; z) - cm(a - 1; c; z) = zm(a; c + 1; z)$
- iii) $(a - 1 + c)m(a; c; z) + (c - a)m(a - 1; c; z) = (c - 1)m(a; c - 1; z)$
- iv) $c(a + z)m(a; c; z) - acm(a + 1; c; z) = (c - a)zm(a; c + 1; z)$

$$v) (c-a)m(a-1; c; z) + (2a-c+z)m(a; c; z) = am(a+1; c; z)$$

$$vi) c(c-1)m(a; c-1; z) - c(c-1+z)m(a; c; z) = (a-c)zm(a; c+1; z)$$

2.4.5 Some example of confluent Hypergeometric function

Example 1

The function

$$erF(z) = \frac{2}{\sqrt{\pi}} \int_0^z \exp(-t^2) dt$$

as defined by (Rainville, 1965) exhibits that

$$erF(z) = \frac{2z}{\sqrt{\pi}} {}_1F_1\left(\frac{1}{2}; \frac{3}{2}; -z^2\right).$$

Solution. Let

$$erF(z) = \frac{2}{\sqrt{\pi}} \int_0^z \exp(-t^2) dt.$$

Then,

$$\begin{aligned} erF(z) &= \frac{2}{\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{(-1)^n \int_0^z t^{2n}}{n!} \\ &= \frac{2}{\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{(-1)^n \int_0^z t^{2n+1}}{n! (2n+1)} \\ &= \frac{2z}{\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{(-1)^n \left(\frac{1}{2}\right)_n z^{2n}}{n! \left(\frac{3}{2}\right)_n} \\ &= \frac{2z}{\sqrt{\pi}} {}_1F_1\left(\frac{1}{2}; \frac{3}{2}; -z^2\right) \end{aligned}$$

Example 2

The incomplete gamma function may be defined by the equation

$$y(a, z) = \int_0^z e^{-t} t^{a-1} dt, R(a) > 0.$$

So that

$$y(a, z) = a^{-1} x^{-a} {}_1F_1(a; a+1; -z).$$

Solution: Let

$$y(a, z) = \int_0^z e^{-t} t^{a-1} dt, R(a) > 0.$$

Then

$$\begin{aligned} y(a, z) &= \int_0^z \sum_{n=0}^{\infty} \frac{(-1)^n t^{n+a-1}}{n!} \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n z^{n+a-1}}{n! (a+n)} \end{aligned}$$

$$\text{now, } (a+n) = \frac{a(a+1)_n}{(a)_n}.$$

Hence

$$y(a, z) = a^{-1} x^a \sum_{n=0}^{\infty} \frac{(-1)^n (a)_n z^n}{n! (a+1)_n} = a^{-1} x^{-a} {}_1F_1(a; a+1; -z).$$

CHAPTER 3

CONFLUENT HYPERGEOMETRIC FUNCTION WITH KUMMER'S FIRST FORMULA

This section introduces the Kummer's first formula and impact with both hypergeometric and confluent hypergeometric function

We can explain the product $e^z \cdot {}_1F_1\left(\begin{smallmatrix} a \\ c \end{smallmatrix}; z\right) = \sum_{n=0}^{\infty} \frac{(-1)^n z^n}{n!} \left(\sum_{k=0}^{\infty} \frac{(a)_k z^k}{(c)_k k!} \right)$

$$= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{(-1)^n z^n}{n!} \frac{(a)_k z^k}{(c)_k k!}$$

When we have

$$\sum_{n=0}^{\infty} \sum_{k=0}^{\infty} A(k, n) = \sum_{n=0}^{\infty} \sum_{k=0}^n A(k, n-k) \quad (3.1)$$

$$= \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{(-1)^{n-k} z^{n-k}}{(n-k)!} \frac{(a)_k z^k}{(c)_k k!}$$

$$\text{and since } (n-k)! = \frac{(-n)_k k!}{(-n)_k}, \quad 0 \leq k \leq n. \quad (3.2)$$

We may write

$$\begin{aligned} &= \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{(-1)^n z^n}{n!} \frac{(-n)_k (a)_k z^k}{(c)_k k!} \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n z^n}{n!} \cdot {}_2F_1(-n, a; c; 1) \end{aligned}$$

But we already know that

$${}_2F_1(-n, a; c; 1) = \frac{\Gamma(c)\Gamma(c-a+n)}{\Gamma(c-a)\Gamma(c+n)}, \frac{\Gamma(c-a+n)}{\Gamma(c-a)} = (b-a)_n, \frac{\Gamma(c+n)}{\Gamma(c)} = (b)_n$$

So that,



$${}_2F_1(-n, a; c; 1) = \frac{(c-a)_n}{(c)_n}$$

then,

$$e^{-z} {}_1F_1(a; c; z) = \sum_{n=0}^{\infty} \frac{(c-a)_n (-z)^n}{(c)_n n!} = {}_1F_1(c-a; c; -z). \quad (3.3)$$

This is Kummer first formula, $c \notin \{-1, -2, -3, \dots, 0\}$

Now under this definition, we will prove the following theorems,

3.1 Theorem

$e^{-t} F(-k, a+n; a; 1) = {}_1F_1(-n; a; -t)$, where k, n are non-negative integer

Proof

$$e^{-t} F(-k, a+n; a; 1) = e^{-t} \sum_{s=0}^k \frac{(-k)_s (a+n)_s}{(a)_s s!}$$

We know that $(a+n)_s = \frac{(a)_{n+s}}{(a)_n}$ (pochhammer property)

So

$$\begin{aligned} e^{-t} F(-k, a+n; a; 1) &= e^{-t} \sum_{s=0}^k \frac{(-k)_s (a)_{n+s}}{s! (a)_s (a)_n} \\ &= \sum_{k=0}^{\infty} \sum_{s=0}^k \frac{(-k)_s (a)_{n+s}}{s! (a)_s (a)_n} \frac{(-1)^k t^k}{k!}, \text{ by (3.2)} \end{aligned}$$

We obtain

$$\begin{aligned} &= \sum_{k=0}^{\infty} \sum_{s=0}^k \frac{(-1)^s k! (a)_{n+s}}{(k-s)! s! (a)_s (a)_n} \frac{(-1)^k t^k}{k!} \\ &= \sum_{k=0}^{\infty} \sum_{s=0}^k \frac{(-1)^s (a)_{n+s}}{s! (a)_s (a)_n} \frac{(-1)^k t^k}{(k-s)!} \end{aligned}$$

Hence

$$\sum_{k=0}^{\infty} \sum_{s=0}^k A(s, k) = \sum_{k=0}^{\infty} \sum_{s=0}^{\infty} A(s, +s) \quad (3.4)$$

So

$$\begin{aligned} \sum_{k=0}^{\infty} \sum_{s=0}^{\infty} \frac{(-1)^s (a)_{n+s} (-1)^{k+s} t^{k+s}}{s! (a)_s (a)_n k!} &= \sum_{k=0}^{\infty} \sum_{s=0}^{\infty} \frac{(a+n)_s t^s (-t)^k}{s! (a)_s k!} \\ &= \sum_{s=0}^{\infty} e^{-t} \frac{(a+n)_s t^s}{(a)_s s!} \\ &= e^{-t} {}_1F_1(+n; a; t) \end{aligned}$$

And since ${}_1F_1(a+n; a; t) = {}_1F_1(-n; a; -t)$ by Kummer's first formula 3.3,

$$\begin{aligned} \sum_{s=0}^{\infty} e^{-t} \frac{(a+n)_s}{(a)_s s!} &= e^{-t} e^t {}_1F_1(-n; a; -t) = {}_1F_1(-n; a; -t) \\ \therefore e^{-t} F(-k, a+n; a; 1) &= {}_1F_1(-n; a; -t) \end{aligned}$$

3.2 Theorem

To prove that

$$\frac{d^k}{dz^k} [{}_1F_1(a+n, a, -t)] = \frac{(-n)_k}{(a)_k} {}_1F_1(a+n, a+k, -t)$$

Proof:

$$\begin{aligned} \frac{d^k}{dz^k} [{}_1F_1(a+n, a, -t)] &= \frac{d^k}{dz^k} [e^{-t} {}_1F_1(-n, a, t)] \quad \text{by (3.3)} \\ &= e^{-t} \left[\frac{d^k}{dz^k} {}_1F_1(-n, a, t) \right] \\ &= e^{-t} \left[\frac{(-n)_k}{(a)_k} {}_1F_1(-n+k, a+k, t) \right] \quad (3.5) \\ &= e^{-t} \left[e^t \frac{(-n)_k}{(a)_k} {}_1F_1(a+k+n-k, a+k, -t) \right] \quad \text{by (3.3)} \end{aligned}$$

$$= \frac{(-n)_k}{(a)_k} {}_1F_1(a+n, a+k, -t)$$

Not that theorem (3.5) is equal to zero if $k=n$

$$\begin{aligned} \frac{d^n}{dz^n} [{}_1F_1(a+n, a, -t)] &= \frac{d^n}{dz^n} [e^{-t} {}_1F_1(-n, a, t)] \text{ by Kummer first formula} \\ &= e^{-t} \left[\frac{d^n}{dz^n} {}_1F_1(-n, a, t) \right] \\ &= e^{-t} \left[\frac{(-n)_n}{(a)_n} {}_1F_1(-n+n, a+n, t) \right] = 0 \end{aligned}$$

Note

Examination of Kummer's first formula soon arouses interest in the special case when the two (CHF) have the same parameters. This happens when $b-a=a$, $b=2a$. we then obtain

$${}_1F_1(a; 2a; z) = e^z {}_1F_1(a; 2a; -z),$$

or

$$e^{-\frac{z}{2}} {}_1F_1(a; 2a; z) = e^{\frac{z}{2}} {}_1F_1(a; 2a; -z) \quad (3.6)$$

More pleasantly, (3.5) may be expressed by saying that the function

$$e^{-z} {}_1F_1(a; 2a; 2z)$$

Is an even function of z . After some step in (Rainville 1967) we get

$$e^z {}_1F_1(a; 2a; -z) = {}_0F_1\left(-; a + \frac{1}{2}; -\frac{1}{4}z^2\right). \quad (3.7)$$

If $2a$ is not an odd integer < 0

Equation (3.7) is known as Kummer's second formula.

3.3 Some example of Kummer's formula with (CHF)

Problem 3.1

Show that

$${}_1F_1(a; b; z) = \frac{1}{\Gamma(a)} \int_0^{\infty} e^{-t} t^{a-1} {}_0F_1(-; b; zt) dt$$

Solution 3.1

We know that

$$\Gamma(z) = \int_0^{\infty} e^{-t} t^{z-1} dt, \quad \operatorname{Re}(z) > 0.$$

Then

$$\begin{aligned} {}_1F_1(a; b; z) &= \sum_{n=0}^{\infty} \frac{(a)_n z^n}{n! (b)_n} \\ &= \frac{1}{\Gamma(a)} \sum_{n=0}^{\infty} \frac{\Gamma(a+n) z^n}{n! (b)_n} \\ &= \frac{1}{\Gamma(a)} \int_0^{\infty} e^{-t} \sum_{n=0}^{\infty} \frac{t^{a+n-1} z^n}{n! (b)_n} dt \\ &= \frac{1}{\Gamma(a)} \int_0^{\infty} e^{-t} t^{a-1} {}_0F_1(-; b; zt) dt, \quad \operatorname{Re}(a) > 0 \end{aligned}$$

Problem 3.2

Show that the aid of the result in problem 1.3, that

$$\int_0^{\infty} \exp(-t^2) t^{2a-n-1} J_n(zt) dt = \frac{\Gamma(a) z^n}{2^{n+1} \Gamma(n+1)} {}_1F_1\left(a; n+1; -\frac{z^2}{4}\right).$$

Solution 3.2

we obtain

$$A = \int_0^{\infty} \exp(-t^2) t^{2a-n-1} J_n(zt) dt = \int_0^{\infty} \frac{e^{-t^2} t^{2a-n-1} z^n t^n}{2^n \Gamma(1+n)} {}_0F_1\left(-; 1+n; -\frac{z^2 t^2}{4}\right) dt$$

Put $t^2 = \beta$. Then

$$A = \frac{z^n}{2^n \Gamma(1+n)} \cdot \frac{1}{2} \int_0^{\infty} e^{-\beta} \beta^{a-1} {}_0F_1\left(-; 1+n; \frac{z^2 \beta^2}{4}\right) d\beta$$

$$= \frac{z^n}{2^{n+1} \Gamma(1+n)} \cdot \frac{\Gamma(a)}{1} {}_1F_1\left(a; n+1; -\frac{z^2}{4}\right)$$

CHAPTER 4

SEVERAL PROPERTIES OF HYPERGEOMETRIC FUNCTION

This chapter proffers an outline of the several properties of hypergeometric function and a detailed discussion of the results.

4.1 Properties

4.1.1 The Contiguous Function relations.

Gauss defined as contiguous to $F(a, b; c; z)$ each of the six function obtained by increasing or decreasing one of the parameters by unity. For simplicity in printing we use the notation,

$$F = F(a, b; c; z)$$

$$F(a+) = F(a+1, b; c; z) \quad (4.1)$$

$$F(a-) = F(a-1, b; c; z) \quad (4.2)$$

Together with similar notations $F(b+)$, $F(b-)$, $F(c+)$ and $F(c-)$ for the other four of the six functions contiguous to F . After some step in (Rainville, 1965) we get this contiguous function relations.

$$i) (a-b)F = aF(a+) - bF(b+)$$

$$ii) (a-c+1)F = aF(a+) - (c-1)F(c-)$$

$$iii) [a + (b-c)z]F = a(1-z)F(a+) - c^{-1}(c-a)(c-b)zF(c+)$$

$$iv) (1-z)F = F(a-) - c^{-1}(c-b)zF(c+)$$

$$v) (1-z)F = F(b-) - c^{-1}(c-a)zF(c+),$$

Example 4.1

From these contiguous functions we can obtain other relations

1) from (iii) and (iv) we get

$$[a + (b - c)z - (c - a)(1 - z)]F = a(1 - z)F(a+) - (c - a)F(a-),$$

In the left hand We get

$$[a + bz - cz - [c - cz - a + az]F = a(1 - z)F(a+) - (c - a)F(a-),$$

So

$$[2a - c + (b - a)z]F = a(1 - z)F(a+) - (c - a)F(a-). \quad (4.3)$$

2) from (iii) and (vi) we get

$$[a + (b - c)z - (c - b)(1 - z)]F = a(1 - z)F(a+) - (c - b)F(b-)$$

So

$$[a + b - c]F = a(1 - z)F(a+) - (c - b)F(b-). \quad (4.4)$$

3) from (2) and (3) we get

$$\begin{aligned} [a + (b - c)z - (a - c + 1)(1 - z)]F \\ = (c - 1)(1 - z)F(c-) - c^{-1}(c - a)(c - b)zF(c+) \end{aligned}$$

Then

$$\begin{aligned} [c - 1 + (a + b - 2c + 1)z]F \\ = (c - 1)(1 - z)F(c-) - c^{-1}(c - a)(c - b)zF(c+). \quad (4.5) \end{aligned}$$

4) from (1) and (4.1) we get

$$\begin{aligned} [(a - b)(1 - z) - 2a + c - (b - a)z]F \\ = (c - a)F(a-) - b(1 - z)F(b+), \end{aligned}$$

Then

$$[c - a - b]F = (c - a)F(a-) - b(1 - z)F(b+). \quad (4.6)$$

4.1.2 Hypergeometric differential equation:

The operator $\theta = z \left(\frac{d}{dz} \right)$, already used in the chapter two of section (2.1.2.3) we resultantly obtained this equation,

$$z(1-z)w'' + [c - (a+b+1)z]w' - abw = 0 \quad (4.7)$$

Example 4.2

In the differential equation (4.7) for $w = F(a, b; c; z)$ introduce a new dependent variable u by $w = (1-z)^{-a}u$, thus obtaining

$$z(1-z)^2u'' + (1-z)[c + (a-b-1)z]u' + a(c-b)u = 0.$$

Next change the independent variable to x by putting $x = \frac{-z}{1-z}$ Show that the equation for u in terms of x is ,

$$x(1-x)\frac{d^2u}{dx^2} + [c - (a+c-b+1)x]\frac{du}{dx} - a(c-b)u = 0, \quad (4.8)$$

And thus derive the solution

$$w = (1-z)^{-1}F\left[\begin{matrix} a, & c-b; \\ c; & \end{matrix} \frac{-z}{1-z}\right]$$

Solution

We know that $w = F(a, b; c; z)$ is a solution of the equation (4.7) in this equation we put $w = (1-z)^{-a}u$ then

$$w' = (1-z)^{-a}u' + a(1-z)^{-a-1}u, \quad (4.9)$$

$$w'' = (1-z)^{-a}u'' + 2a(1-z)^{-a-1}u' + a(a+1)(1-z)^{-a-2}u. \quad (4.10)$$

Now we get the new equation from the eq (4.8),(4.9)and (4.7)

$$\begin{aligned} z(1-z)u'' + 2azu' + a(a+1)z(1-z)^{-1}u + cu' + ca(1-z)^{-1}u - (a+b+1)zu' \\ - a(a+b+1)z(1-z)^{-1}u - abu = 0, \end{aligned}$$

Then

$$z(1-z)^2 + (1-z)[c + (a-b-1)z]u' + a(c-b)u = 0. \quad (4.11)$$

Now put $x = \frac{-z}{1-z}$, then $z = \frac{-x}{1-x}$, $x = \frac{-z}{1-z}$, $1-z = \frac{1}{1-x}$ so $\frac{dx}{dz} = \frac{-1}{(1-z)^2} = -(1-x)^2$

$$\frac{d^2x}{dz^2} = \frac{-2}{(1-z)^3} = -2(1-x)^3$$

The old equation (4.11) above may be written

$$\frac{d^2u}{dz^2} + \left[\frac{c}{z(1-z)} + \frac{a-b-1}{1-z} \right] \frac{du}{dz} + \frac{a(c-b)}{z(1-z)^2} u = 0,$$

Which then leads to the new equation

$$(1-x)^4 \frac{d^2u}{dx^2} + \left[-2(1-x)^3 - (1-x)^2 \left\{ \frac{c(1-x)^2}{-x} + (a-b-1)(1-x) \right\} \right] \frac{du}{dx} - \frac{a(c-b)(1-x)^3}{x} u = 0$$

or

$$x(1-x) \frac{d^2u}{dx^2} + [-2x - \{-c(1-x) + (a-b-1)x\}] \frac{du}{dx} - a(c-b)u = 0,$$

Or

$$x(1-x) \frac{d^2u}{dx^2} + [x - (a-b+c+1)x] \frac{du}{dx} - a(c-b)u = 0 \quad (4.12)$$

Now (4.12) is a hypergeometric equation with parameters $\gamma = c, \alpha + \beta + 1$

$a-b+c+1, \alpha\beta = a(c-b)$. Hence $\alpha = a, \beta = c-b, \gamma = c$. One solution of (4.12) is

$$u = F(a, c-b; c; x),$$

So one solution of equation (4.7) is

$$w = (1-z)^{-1} F \left[\begin{matrix} a, c-b; \\ c; \end{matrix} \frac{-z}{1-z} \right]$$

4.1.3 Elementary series manipulation

(Choi, 2003) established some generalized principles of double series manipulations some special cases of which are also written for easy reference in their use. Note that $A_{x,y}$ denotes a function of two variables x and y , and N is the set of positive integers

$$1) \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} A_{k;n} = \sum_{n=0}^{\infty} \sum_{k=0}^n A_{k;n-k};$$

$$2) \sum_{n=0}^{\infty} \sum_{k=0}^n A_{k;n} = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} A_{k;n+k};$$

$$3) \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} A_{k;n} = \sum_{n=0}^{\infty} \sum_{k=0}^{\frac{n}{2}} A_{k;n-2k}$$

$$4) \sum_{n=0}^{\infty} \sum_{k=0}^{\frac{n}{2}} A_{k;n} = \sum_{n=0}^{\infty} \sum_{k=0}^n A_{k;n+2k}$$

Example 4.2

Prove that if $g_n = F(-n, \alpha; 1 + \alpha - n; 1)$ and α is not an integer, then $g_n = 0$ for $n \geq 1$, $g_0 = 1$.

Solution

Let $g_n = F(-n, \alpha; 1 + \alpha - n; 1)$.

Then

$$g_n = \sum_{k=0}^n \frac{(-n)_k (a)_k}{k! (1 + a - n)_k} = \sum_{k=0}^n \frac{n! (-a)_k (a)_k}{n! (k-1)! (a)_n}$$

Hence compute the series

$$\begin{aligned}
\sum_{n=0}^{\infty} \frac{(-a)_n g_n t^n}{n!} &= \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{(a)_k (-a)_{n-k} t^n}{k! (n-k)!} \\
&= \left(\sum_{n=0}^{\infty} \frac{(a)_n t^n}{n!} \right) \left(\sum_{n=0}^{\infty} \frac{(-a)_n t^n}{n!} \right) \\
&= (1-t)^a (1-t)^{-a} \\
&= 1
\end{aligned}$$

Therefore, $g_0 = 1$ and $g_n = 0$ for $n \geq 1$. (Note: easiest to choose $\alpha \neq$ integer, can actually do better than that probably).

4.1.5 A quadratic transformation

A quadratic transformation as established by (Rainville, 1965) is based on the following;

4.1.5.1 Theorem

If $2b$ is neither zero nor negative integer and if both $|x| < 1$ and $|4x(1+x)^{-2}| < 1$

$$(1+x)^{-2a} F \left[\begin{matrix} a, b; \\ 2b; \end{matrix} \frac{4x}{(1+x)^2} \right] = F \left[\begin{matrix} a, a-b+\frac{1}{2}; \\ b+\frac{1}{2}; \end{matrix} x^2 \right].$$

Example 4.5

In this theorem put $b = \alpha, a = \alpha + \frac{1}{2}$, $4x(1+x)^{-2} = z$ and thus prove that

$$\left[\begin{matrix} \alpha, \alpha + \frac{1}{2}; \\ 2\alpha; \end{matrix} z \right] = (1-z)^{\frac{1}{2}} \left[\frac{2}{1+\sqrt{1-z}} \right]^{2\alpha-1}$$

Solution

Theorem 2 gives us

$$(1+x)^{-2a} F \left[\begin{matrix} a, b; \\ ab; \end{matrix} \frac{4x}{(1+x)^2} \right] = F \left[\begin{matrix} a, a-b+\frac{1}{2}; \\ b+\frac{1}{2}; \end{matrix} x^2 \right].$$

Then put $b = \alpha, a = \alpha + \frac{1}{2}, 4x(1+x)^{-2} = z$

then

$$zx^2 + 2(z-2)x + z = 0$$

$$zx = 2 - 3 \pm \sqrt{z^2 - 4z + 4 - 3^2} = 2 - z \pm 2\sqrt{1-z}$$

Now $x = 0$ when $z = 0$, so

$$zx = 2 - z - 2\sqrt{1-z} = 1 - z + 1 - 2\sqrt{1-z}$$

Therefore

$$x = \frac{(1 - \sqrt{1-z})^2}{z} = \frac{(1 - \sqrt{1-z})[1 - (1-z)]}{z(1 + \sqrt{1-z})}.$$

Thus

$$x = \frac{1 - \sqrt{1-z}}{1 + \sqrt{1-z}}$$

And

$$x + 1 = \frac{2}{(1 + \sqrt{1-z})}$$

Then we obtain

$$\frac{4x}{(1+x)^2} = \frac{(1 - \sqrt{1-z})}{(1 + \sqrt{1-z})} \cdot \frac{(1 - \sqrt{1-z})^2}{4} = z$$

a check. Now with $b = \alpha, a = \alpha + 1$ theorem 4 yields

$$\left[\frac{2}{1 + \sqrt{1-z}} \right]^{2\alpha-1} F \left[\begin{matrix} \alpha + \frac{1}{2}, \alpha; \\ 2\alpha; \end{matrix} z \right] = \left[\begin{matrix} \alpha + \frac{1}{2}, 1; \\ \alpha + \frac{1}{2}; \end{matrix} x^2 \right] = F \left[\begin{matrix} 1; \\ -; \end{matrix} x^2 \right] = (1-x^2)^{-1}.$$

Since

$$1-x = \frac{1 - \sqrt{1-z}}{1 + \sqrt{1-z}} \text{ and } 1+x = \frac{2}{1 + \sqrt{1-z}}.$$

$$(1-x^2) = \frac{4\sqrt{1-z}}{(1 + \sqrt{1-z})^2}$$

Thus we have

$$\begin{aligned} F \left[\begin{matrix} \alpha, \alpha + \frac{1}{2}; \\ 2\alpha; \end{matrix} z \right] &= \left[\frac{2}{1 + \sqrt{1-z}} \right]^{2\alpha+1} \cdot \left[\frac{2}{1 + \sqrt{1-z}} \right]^{-2} (1-z)^{-\frac{1}{2}} \\ &= (1-z)^{\frac{1}{2}} \left[\frac{2}{1 + \sqrt{1-z}} \right]^{2\alpha-1}, \end{aligned}$$

as defined. Now we use theorem 3 to see that

$$F \left[\begin{matrix} \alpha, \alpha + \frac{1}{2}; \\ 2\alpha; \end{matrix} z \right] = (1-z)^{-\frac{1}{2}} F \left[\begin{matrix} \alpha, \alpha - \frac{1}{2}; \\ 2\alpha; \end{matrix} z \right]$$

So that we also get

$$F \left[\begin{matrix} \alpha, \alpha - \frac{1}{2}; \\ 2\alpha; \end{matrix} z \right] = \left[\frac{2}{1 + \sqrt{1-z}} \right]^{2\alpha-1},$$

as desired.

4.1.6 Additional properties

We will obtain one more identity as an example of those resulting from combination of the theorem proved earlier in this chapter. In the Identity of theorem 3, replace a by $(\frac{1}{2}c - \frac{1}{2}a)$ and b by $(\frac{1}{2}c + \frac{1}{2}a - \frac{1}{2})$ to get

$$F\left[\begin{matrix} \frac{1}{2}c - \frac{1}{2}a, \frac{1}{2}c + \frac{1}{2}a - \frac{1}{2} \\ c \end{matrix}; 4x(1-x)\right] = F\left[\begin{matrix} c-a, c+a-1 \\ c \end{matrix}; x\right].$$

Theorem 1 yields

$$F\left[\begin{matrix} c-a, c+a-1 \\ c \end{matrix}; x\right] = (1-x)^{1-c} F\left[\begin{matrix} a, 1-a \\ c \end{matrix}; x\right],$$

Which leads to the desired result.

4.1.6.1 Theorem

If c is neither zero nor negative integer and if both $|x| < 1$ and $|4x(1-x)| < 1$

$$F\left[\begin{matrix} a, 1-a \\ c \end{matrix}; x\right] = (1-x)^{1-c} F\left[\begin{matrix} \frac{1}{2}c - \frac{1}{2}a, \frac{1}{2}c + \frac{1}{2}a - \frac{1}{2} \\ c \end{matrix}; 4x(1-x)\right].$$

Example

Use this theorem to show that

$$(1-x)^{1-c} F\left[\begin{matrix} a, 1-a \\ c \end{matrix}; x\right] = (1-2x)^{a-c} F\left[\begin{matrix} \frac{1}{2}c - \frac{1}{2}a, \frac{1}{2}c - \frac{1}{2}a + \frac{1}{2} \\ c \end{matrix}; \frac{4x(1-x)}{(1-2x)^2}\right]$$

Solution

$$(1-x)^{1-c} F\left[\begin{matrix} a, 1-a \\ c \end{matrix}; x\right] = F\left[\begin{matrix} \frac{c-a}{2}, \frac{c+a-1}{2} \\ c \end{matrix}; 4x(1-x)\right]$$

CHAPTER 5

CONCLUSION AND SUGGESTIONS FOR FUTURE

This study had presented definitions and examples of hypergeometric function; confluent hypergeometric function, and Kummer confluent hypergeometric function. It can therefore be concluded that theorems and some properties. Moreover, it can also be concluded that the Kummer function has wide application in various subjects and hence proving stability or other properties were drawn to be of paramount importance. This study centered on the Kummer's first formula with confluent hypergeometric function. Future studies can endeavor to extend insights on this area in depth.

REFERENCES

- Andrews, L. C. (1998). Special functions of mathematics for engineers. *A publication of SPIE-the International Society for Optical Engineering Bellingham, Washington USA*, 49, 194-261.
- Barnes, E. W. (1908). A new development of the theory of the hypergeometric functions. In *Proceedings of the London Mathematical Society* (pp. 141-177). London: Cambridge University.
- Buchholz, H. (2013). The confluent hypergeometric function: with special emphasis on its applications. *Springer Science & Business Media*, 15, 34-73.
- Campos, L. M. B. C. (2001). On some solutions of the extended confluent hypergeometric differential equation. *Journal of Computational and Applied Mathematics*, 137(1), 177- 200.
- Choi, J. (2003). Notes on formal manipulations of double series. *Communications Korean mathematical society*, 18(4), 781-789.
- Dickenstein, A. (2004). Hypergeometric functions with integer homogeneities. Retrieved June 29, 2016 from: <http://mate.dm.uba.ar/~alidick/papers/paris03.pdf>
- die reine und angewandte Mathematik*, 17, 228-242.
- Freden, W., & Gutting, M. (2013). Special functions of mathematical (geo-) physics. Springer Science & Business Media. New York: Addison Wesley.
- Georgiev, G. N., & Georgieva-Grosse, M. N. (2005). The Kummer confluent hypergeometric function and some of its applications in the theory of azimuthally magnetized circular ferrite waveguides. *Journal of Telecommunications and Information Technology*, 112-128.
- Gogolin, A. O. (2013). Lectures on Complex Integration. Springer Science & Business Media. New York: Addison Wesley.
- Gronau, D. (2003). Why is the gamma function so as it is. *Journal of Teaching Mathematics and Computer Science*, 1(7), 43-53. Ko, M. S. (2011). Approximation and analysis of

- confluent hypergeometric differential equation in homing guidance. Retrived June 29, 2016 from: http://amath.kaist.ac.kr/pde_lab/thesis_papers/Master1102MinSuKo.pdf
- Kummer, E. E. (1837). De integralibus quibusdam definitis et seriebus infinitis. *Journal für Die Reine und Angewandte Mathematik*, 17, 228-242.
- Luke, Y. L. (1972). Inequalities for generalized hypergeometric functions. *Journal of Approximation Theory*, 5(1), 41-65.
- Özergin, E. (2011). Some properties of hypergeometric functions. Retrived June 29, 2016 from: <http://irep.emu.edu.tr:8080/xmlui/bitstream/handle/11129/217/Ozergin.pdf?sequence=1>
- Pearson, J. (2009). Computation of hypergeometric functions. Retrieved June 29, 2016 from: https://www.researchgate.net/profile/John_Pearson16/publication/265047721_Computation_of_Hypergeometric_Functions/links/55d5f38408aec156b9a6fafd.pdf
- Rainville, E. D. (1960). Special mathematical functions. Macmillan: New York, NY.
- Sebah, P., & Gourdon, X. (2002). Introduction to the gamma function numbers, computation free and constants. Retrieved June 29, 2016 from: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.527.8519&rep=rep1&type=pdf>

[Ödevler](#)[Öğrenciler](#)[Kütüphaneler](#)[Takvim](#)[Tartışma](#)[Tercihler](#)

Bu sayfa hakkında

Bu sizin ödev gelen kutunuzdur. Bir ödevi görüntülemek için, ödev başlığına tıklayın. Orijinallik Raporu'nu görmek için, benzerlik raporundaki orijinallik raporu ikonuna tıklayın. Bu ikon tıklanabilir durumda değilse, orijinallik raporu henüz oluşturulmamış demektir.

SHWAN SWARA FATAH

Gelen Kutusu | Görüntüleniyor: yeni ödevler ▼

Dosyayı Gönder Ödev ayarlarını düzenle | E-posta bildirmeyenler

<input type="checkbox"/>	Yazar	Başlık	Benzerlik	web	yayın	student papers	Dosya	Ödev Numarası	Tarih
<input type="checkbox"/>	Shwan Fatah	CONCLUSION	%0 <input type="text" value="%0"/>	0%	0%	0%	ödev indir	687371790	01-Tem-2016
<input type="checkbox"/>	Shwan Swara Fatah	ABSTACT	%0 <input type="text" value="%0"/>	0%	0%	0%	ödev indir	685933094	23-Haz-2016
<input type="checkbox"/>	Shwan Swara Fatah	CHAPTER 3	%4 <input type="text" value="%4"/>	3%	2%	0%	ödev indir	685933215	23-Haz-2016
<input type="checkbox"/>	Shwan Swara Fatah	CHAPTER 2	%12 <input type="text" value="%12"/>	4%	7%	2%	ödev indir	685933174	23-Haz-2016
<input type="checkbox"/>	Shwan Swara Fatah	ALL DOC.	%16 <input type="text" value="%16"/>	14%	4%	4%	ödev indir	685933623	23-Haz-2016
<input type="checkbox"/>	Shwan Swara Fatah	CHAPTER 1	%17 <input type="text" value="%17"/>	14%	8%	5%	ödev indir	685933139	23-Haz-2016
<input type="checkbox"/>	Shwan Swara Fatah	CHAPTER 4	%17 <input type="text" value="%17"/>	12%	10%	2%	ödev indir	685933262	23-Haz-2016

